

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

December 06, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

**APPLICATION NUMBER: 10/635,795
FILING DATE: August 06, 2003
RELATED PCT APPLICATION NUMBER: PCT/US04/25395**

Certified by

Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office



BEST AVAILABLE COPY

Please type a plus sign (+) inside this box → +

PTO/SB/05 (03-01)

Approved for use through 10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL		Attorney Docket No. 010023-000800US	
		First Inventor	John H. Crowe, et al.
		Title	CELLS AND METHOD FOR PRESERVING CELLS
(Only for new nonprovisional applications under 37 C.F.R. 1.53(b))		Express Mail Label No.	Eu121998655US


APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents.</i> 1. <input checked="" type="checkbox"/> Fee Transmittal Form (e.g., PTO/SB/17) <i>(Submit an original and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. 3. <input checked="" type="checkbox"/> Specification [Total Pages 62] <i>(preferred arrangement set forth below)</i> <ul style="list-style-type: none">- Descriptive title of the invention- Cross Reference to Related Applications- Statement Regarding Fed sponsored R & D- Reference to sequence listing, a table, or a computer program listing appendix- Background of the invention- Brief Summary of the invention- Brief Description of the Drawings (if filed)- Detailed Description- Claim(s)- Abstract of the Disclosure 4. <input checked="" type="checkbox"/> Drawing(s) (35 U.S.C.113) [Total Sheets 21] 5. Oath or Declaration [Total Pages 5] <ul style="list-style-type: none">a. <input type="checkbox"/> Newly executed (original or copy)b. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63 (d)) <i>(for a continuation/divisional with Box 18 completed)</i>i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 6. <input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76	ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231 7. <input type="checkbox"/> CD-ROM or CD-R in duplicate, large table or Computer Program <i>(Appendix)</i> 8. Nucleotide and/or Amino Acid Sequence Submission <i>(if applicable, all necessary)</i> <ul style="list-style-type: none">a. <input type="checkbox"/> Computer Readable Form (CRF)b. Specification Sequence Listing on:<ul style="list-style-type: none">i. <input type="checkbox"/> CD-ROM or CD-R (2 copies); orii. <input type="checkbox"/> paperc. <input type="checkbox"/> Statements verifying identity of above copies ACCOMPANYING APPLICATIONS PARTS 9. <input type="checkbox"/> Assignment Papers (cover sheet & document(s)) 10. <input type="checkbox"/> 37 C.F.R. §3.73(b) Statement <input type="checkbox"/> Power of Attorney <i>(when there is an assignee)</i> 11. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 12. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations 13. <input type="checkbox"/> Preliminary Amendment 14. <input type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i> 15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i> 16. <input type="checkbox"/> Non publication Request under 35 U.S.C. 122 (b)(2)(B)(i). Applicant must attach form PTO/SB/35 or its equivalent. 17. <input checked="" type="checkbox"/> Other: <u>Express Mail Certification</u>
--	---

18. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment, or in an Application Data Sheet under 37 CFR 1.76:

<input type="checkbox"/> Continuation	<input type="checkbox"/> Divisional	<input type="checkbox"/> Continuation-in-part (CIP)	of prior application No: _____ / _____
Prior application information: Examiner _____			Group / Art Unit: _____

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 5b, is considered a part of the disclosure of the accompanying or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number or Bar Code Label		37490		or <input type="checkbox"/> Correspondence address below	
Name	John W. Carpenter				
Address	Carpenter & Kulas, LLP 1900 Embarcadero Road, Suite 109				
City	Palo Alto	State	CA	Zip Code	94303
Country	USA	Telephone	650-842-0303	Fax	650-842-0304

Name (Print/Type)	John W. Carpenter	Registration No. (Attorney/Agent)	26,447
Signature		Date	August 6, 2003

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

10/635795
08/06/03

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**FEE TRANSMITTAL
for FY 2002**

Patent fees are subject to annual revision.

Complete if Known

Application Number	Unknown
Filing Date	August 6, 2003
First Named Inventor	John H. Crowe, et al.
Examiner Name	Unknown
Group / Art Unit	Unknown
Attorney Docket No.	010023-000800US

TOTAL AMOUNT OF PAYMENT (\$) 856.00**METHOD OF PAYMENT (check one)**

- 1.
- ☐
- The Commissioner is hereby authorized to charge indicated fees and credit any over payments to:

Deposit
Account
NumberDeposit
Account
Name☐ Charge Any Additional Fee Required
Under 37 CFR 1.16 and 1.17☒ Applicant claims small entity status.
See 37 CFR 1.27

- 2.
- ☒
- Payment Enclosed:

☒ Check ☐ Credit card ☐ Money
Order ☐ Other**FEE CALCULATION****1. BASIC FILING FEE**

Large Fee Code	Entity Fee (\$)	Small Fee Code	Entity Fee (\$)	Fee Description	Fee Paid
101	740	201	370	Utility filing fee	370.00
106	330	206	165	Design filing fee	
107	510	207	255	Plant filing fee	
108	740	208	370	Reissue filing fee	
114	160	214	80	Provisional filing fee	

SUBTOTAL (1)

(\$ 370.00)

2. EXTRA CLAIM FEES

Total Claims	Independent Claims	Multiple Dependent	Extra Claims	Fee from below	Fee Paid
33	-20		13	9	117
6	-3		3	42	126
					0

Large Fee Code	Entity Fee (\$)	Small Fee Code	Entity Fee (\$)	Fee Description
103	18	203	9	Claims in excess of 20
102	84	202	42	Independent claims in excess of 3
104	280	204	140	Multiple dependent claim, if not paid
109	84	209	42	** Reissue independent claims over original patent
110	18	210	9	** Reissue claims in excess of 20 and over original patent

SUBTOTAL (2)

(\$ 243.00)

**or number previously paid, if greater; For Reissues, see above

FEE CALCULATION (continued)

Fee Code	Large Entity Fee (\$)	Small Entity Fee Code	Small Entity Fee (\$)	Fee Description	Fee Paid
105	130	205	65	Surcharge - late filing fee or oath	
127	50	227	25	Surcharge - late provisional filing fee or cover sheet	
139	130	139	130	Non-English specification	
147	2,520	147	2,520	For filing a request for reexamination	
112	920*	112	920*	Requesting publication of SIR prior to Examiner action	
113	1,840*	113	1,840*	Requesting publication of SIR after Examiner action	
115	110	215	55	Extension for reply within first month	
116	400	216	200	Extension for reply within second month	
117	920	217	460	Extension for reply within third month	
118	1,440	218	720	Extension for reply within fourth month	
128	1,960	228	980	Extension for reply within fifth month	
119	320	219	160	Notice of Appeal	
120	320	220	160	Filing a brief in support of an appeal	
121	280	221	140	Request for oral hearing	
138	1,510	138	1,510	Petition to institute a public use proceeding	
140	110	240	55	Petition to revive - unavoidable	
141	1,280	241	640	Petition to revive - unintentional	
142	1,280	242	640	Utility issue fee (or reissue)	
143	460	243	230	Design issue fee	
144	620	244	310	Plant issue fee	
122	130	122	130	Petitions to the Commissioner	
123	50	123	50	Petitions related to provisional applications	
126	180	126	180	Submission of Information Disclosure Stmt	
581	40	581	40	Recording each patent assignment per property (times number of properties)	
146	740	246	370	Filing a submission after final rejection (37 CFR § 1.129(a))	
149	740	249	370	For each additional invention to be examined (37 CFR § 1.129(b))	
179	740	279	370	Request for Continued Examination (RCE)	
169	900	169	900	Request for expedited examination of a design application	


Other fee (specify) _____

*Reduced by Basic Filing Fee Paid

SUBTOTAL (3)

(\$)

SUBMITTED BY**Complete (if applicable)**

Name (Print/Type)	John W. Carpenter	Registration No. Attorney/Agent	26,447	Telephone	650-842-0303
Signature				Date	August 6, 2003

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

In Re Application of: John H. Crowe, et al. Application No.: Unknown Filing Date: August 6, 2003 Title: CELLS AND METHOD FOR PRESERVING CELLS	Docket No.: 010023-000800US Re: Filing Patent Application
---	--

Commissioner for Patents
 P.O. Box 1450
 Alexandria, VA 22313-1450

EXPRESS MAIL CERTIFICATE

Utility Transmittal Letter;
 Fee Transmittal Sheet;
 Unsigned Declaration
 Specification, Claims, Abstract and Drawings


I hereby certify that this paper and the enclosures listed above are being deposited with the U.S. Postal Service "Express Mail to Addressee" Express Mail No.: EU121998655US on August 6, 2003, and is addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Dated: August 6, 2003

Respectfully submitted,

EU 121998655 US

The Regents of the University of California

By: 
 John W. Carpenter, Reg. No. 26,447
 CARPENTER & KULAS, LLP
 1900 Embarcadero Road
 Suite 109
 Palo Alto, CA 94303
 USA

UC Davis Case No. 2003-337-1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION FOR PATENT

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe
Fern Tablin
Nelly Tsvetkova
Zsolt Torok
Rachna Bali
Gyana Satpathy
Denis Dwyre

Related Patent Applications

This patent application is related to co-pending patent application Serial No. 10/052,162, filed January 16, 2002. Patent application Serial No. 10/052,162 is a continuation-in-part patent application of co-pending patent application Serial No. 09/927,760, filed August 9, 2001. Patent application Serial No. 09/927,760 is a continuation-in-part patent application of co-pending patent application Serial No. 09/828,627, filed April 5, 2001. Patent application Serial No. 09/828,627 is a continuation patent application of patent application Serial No.

09/501,773, filed February 10, 2000. All of the foregoing patent applications are fully incorporated herein by reference thereto as if repeated verbatim immediately hereinafter.

Field of the Invention

Embodiments of the present invention generally broadly relate to living mammalian cells. More specifically, embodiments of the present invention generally provide for the preservation and survival of cells, especially human cells, such as erythrocytic cells.

Embodiments of the present invention also generally broadly relate to the therapeutic uses of cells; and more particularly to manipulations or modifications of erythrocytic cells, such as loading erythrocytic cells with solutes and in preparing freeze-dried compositions that can be re-hydrated at the time of application. When cells for various embodiments of the present invention are re-hydrated, they are immediately restored to viability.

The compositions and methods for embodiments of the present invention are useful in many applications, such as in medicine, pharmaceuticals, biotechnology, and agriculture, and including transfusion therapy, as hemostasis aids and for drug delivery.

Statement Regarding Federal Sponsored Research and Development

Embodiments of this invention were made with Government support under Grant No. N66001-00-C-8048, awarded by the Department of Defense Advanced Research Projects Agency (DARPA). Further embodiments of this invention were made with Government support under Grant Nos. HL57810 and HL61204, awarded by the

National Institutes of Health. The Government has certain rights to embodiments of this invention.

Background of the Invention

A cell is broadly regarded in the art as a small, typically microscopic, mass of protoplasm bounded externally by a semi-permeable membrane, usually including one or more nuclei and various other organelles with their products. A cell is capable either alone or interacting with other cells of performing all the fundamental function(s) of life, and forming the smallest structural unit of living matter capable of functioning independently.

Cells may be transported and transplanted; however, this requires cryopreservation which includes freezing and subsequent reconstitution (e.g., thawing, re-hydration, etc.) after transportation. Unfortunately, a very low percentage of cells retain their functionality after undergoing freezing and thawing. While some cryoprotectants, such as dimethyl sulfoxide, tend to lessen the damage to cells, they still do not prevent some loss of cell functionality.

Trehalose has been found to be suitable in the cryopreservation of cells and platelets. Trehalose is a disaccharide found at high concentrations in a wide variety of organisms that are capable of surviving almost complete dehydration. Trehalose has been shown to stabilize membranes, proteins, and certain cells during freezing and drying in vitro.

U.S. Patent No. 5,827,741, Beattie et al., issued October 27, 1998, discloses cryoprotectants for human cells and platelets, such as dimethylsulfoxide and trehalose. The cells or platelets may be suspended, for example, in a solution

containing a cryoprotectant at a temperature of about 22°C and then cooled to below 15°C. This incorporates some cryoprotectant into the cells or platelets, but not enough to prevent hemolysis of a large percentage of the cells or platelets.

Accordingly, a need exists for the effective and efficient preservation of cells. More specifically, and accordingly further, a need also exists for the effective and efficient cryopreservation of cells (e.g., erythrocytic cells, eukaryotic cells, or any other cells, and the like), such that the preserved cells respectively maintain their biological properties and may readily become viable after storage.

Summary of Embodiments of the Invention

In one aspect of the present invention, a dehydrated composition is provided having a generally dehydrated composition comprising freeze-dried cells selected from a mammalian species (e.g., a human) and being effectively loaded internally (e.g., producing hyper-osmotic pressure on the cells to uptake external trehalose via fluid phase endocytosis) with at least about 10 mM of a carbohydrate (e.g., an oligosaccharide, such as trehalose) therein to preserve biological properties during freeze-drying and re-hydration. The amount of the carbohydrate inside the freeze-dried cells is preferably the amount obtained from maintaining a positive loading gradient or loading efficiency gradient on the cell. When the carbohydrate is trehalose, the amount of trehalose loaded inside the freeze-dried cells is preferably from about 10 mM to about 50 mM.

In another aspect of the present invention, a method is provided for loading (e.g., by fluid phase endocytosis) a solute into a cell (e.g., an erythrocytic cell). Embodiments of the invention include disposing a cell in a solution having a solute concentration of sufficient magnitude to produce hyper-osmotic pressure on the cell for transferring a solute (e.g., an oligosaccharide, such as trehalose) from the solution into the cell. The method may additionally comprise preventing a decrease in a loading efficiency gradient in the loading of the solute into the cell. In an embodiment of the invention where the solute comprises an oligosaccharide, the preventing a decrease in a loading efficiency gradient in the loading of the oligosaccharide into the cell may comprise maintaining a concentration of the oligosaccharide in the oligosaccharide solution below a certain concentration, such as below from about 35 mM to about 65 mM, more particularly below a concentration ranging from about 40 mM to about 60 mM, more particularly further below a concentration ranging from about 45 mM to about 55 mM (e.g., below about 50 mM). In another embodiment of the invention, the preventing a decrease in a loading efficiency gradient in the loading of the oligosaccharide into the cell comprises maintaining a positive gradient of loading efficiency to concentration of the oligosaccharide in the oligosaccharide solution.

The solute concentration includes an extracellular cellular solute concentration for elevating extracellular osmolarity within the solution to a value which is greater than a value of the intracellular osmolarity of the cell. The transferring of the solute is preferably by fluid phase endocytosis and preferably without degradation of the solute. In embodiments of the invention where the cell is an erythrocytic cell and the solute comprises trehalose, a gradient of trehalose (mM) within

the erythrocytic cell to extracellular trehalose concentration (mM) within the solution may range from about 0.130 to about 0.200, particularly for a temperature ranging from about 30° C to about 40° C (e.g., about 37° C). In a further embodiment of the invention, a gradient of trehalose (mM) within the erythrocytic cell to extracellular trehalose concentration (mM) within the solution ranges from about 0.04 to about 0.12, particularly for a temperature ranging from about 0° C to about 10° C. In yet a further embodiment, a gradient of trehalose (mM) within the erythrocytic cell to extracellular trehalose concentration (mM) within the solution may range from about 0.04 to about 0.08, or from about 0.08 to about 0.12, particularly for a temperature ranging from about 0° C to about 10° C. The solute solution may have a trehalose concentration ranging from about 320 mM to about 4000 mM, such as including from about 320 mM to about 2000 mM or from about 500 mM to about 1000 mM.

A further embodiment of the invention provides retaining the solute in the cell; more specifically, washing the cell and retaining the solute in the cell during the washing. The washing is with a washing buffer, and retention of the solute in the cell increases from about 25% to about 175% when a buffer concentration (e.g., the osmolarity of all osmotically active particles within the washing buffer solution) increases from about 50% to about 400%, more preferably from about 50% to about 150% when a buffer concentration increases from about 100% to about 300%, and most preferably from about 75% to about 125% (e.g., about 100%) when a buffer concentration increases from about 150% to about 250% (e.g., about 200%). The washing of the cell with a washing buffer includes employing a ratio of an extracellular buffer concentration (mOsm) to an intracellular solute concentration (mM) ranging from about 14.0 to about 4.0,

such as from about 12.0 to about 5.0, including from about 9.0 to about 6.0 and from about 8.0 to about 7.0 (e.g., about 7.5).

Additional embodiments of the present invention provide a method for loading trehalose into an erythrocytic cell. The method may comprise disposing an erythrocytic cell in a trehalose solution having a trehalose concentration of at least about 25 % (preferably at least about 50%) greater than the intracellular osmolarity of the erythrocytic cell for loading (e.g., by fluid phase endocytosis) the trehalose into the erythrocytic cell.

The loading of the trehalose from the trehalose solution into the erythrocytic cell may be without degradation of the trehalose, and produces a loaded erythrocytic cell having a gradient of loaded trehalose (mM) within the erythrocytic cell to extracellular trehalose concentration (mM) within the trehalose solution ranging from about 0.130 to about 0.200. In another embodiment, the loading of the trehalose produces a loaded erythrocytic cell having a gradient of loaded trehalose (mM) within the erythrocytic cell to extracellular trehalose concentration (mM) within the trehalose solution ranging from about 0.04 to about 0.12. In a further embodiment, the loading of the trehalose produces a loaded erythrocytic cell having a gradient of loaded trehalose (mM) within the erythrocytic cell to extracellular trehalose concentration (mM) within the trehalose solution ranging from about 0.04 to about 0.08, or from about 0.08 to about 0.12, depending on the extracellular trehalose concentration and the temperature of the trehalose solution. The trehalose solution may have a trehalose concentration ranging from about 25 % to at least about 1000 % greater than the intracellular osmolarity of the erythrocytic cell, or at least about 50% greater than the intracellular osmolarity of the erythrocytic cell.

A further embodiment of the invention provides retaining the trehalose in the erythrocytic cell; more specifically washing the erythrocytic cell and retaining the trehalose in the erythrocytic cell during the washing.

The washing of the erythrocytic cell is preferably with a washing buffer, and retention of the trehalose in the erythrocytic cell increases from about 25% to about 175% when a buffer concentration increases from about 50% to about 400%, more preferably from about 50% to about 150% when a buffer concentration increases from about 100% to about 300%, and most preferably from about 75% to about 125% (e.g., about 100%) when a buffer concentration increases from about 150% to about 250% (e.g., about 200%). The washing of the erythrocytic cell with a washing buffer includes employing a ratio of an extracellular buffer concentration (mOsm) to an intracellular trehalose concentration (mM) ranging from about 14.0 to about 4.0, more particularly from about 12.0 to about 5.0, including from about 9.0 to about 6.0 and from about 8.0 to about 7.0 (e.g., about 7.5).

Additional embodiments of the present invention provide a method for loading (e.g., by fluid phase endocytosis) an oligosaccharide into cells (e.g., erythrocytic cells) comprising disposing cells in an oligosaccharide solution having an oligosaccharide concentration of at least about 25 % greater than the intracellular osmolarity of the cells for loading oligosaccharide into the cells, and preventing a decrease in a loading gradient in the loading of the oligosaccharide into the cells. In one embodiment of the invention, the preventing a decrease in a loading gradient in the loading of the oligosaccharide into the cells comprises maintaining a concentration of the oligosaccharide in the oligosaccharide solution below a certain concentration, such as below a

concentration ranging from about 35 mM to about 65 mM, more particularly below a concentration ranging from about 40 mM to about 60 mM, more particularly further below a concentration ranging from about 45 mM to about 55 mM (e.g., below about 50 mM). In another embodiment the preventing a decrease in a loading gradient in the loading of the oligosaccharide into the cells comprises maintaining a positive gradient of concentration of oligosaccharide loaded into the cells to concentration of the oligosaccharide in the oligosaccharide solution.

Further embodiments of the present invention provide a method for preparing a dehydrated composition comprising loading cells in a loading solution having a salt solution and a solute for producing loaded cells, and lyophilizing the loaded cells in a freeze-drying solution having a drying-salt solution, the solute, an inert substance and a protein to produce a dehydrated composition. The loading solution may comprise at least about 200 mM of the solute and at least about 75 mOsm of the salt solution. The freeze-drying solution may comprise at least about 50 mM of the solute, at least about 2.0 % by weight of the inert substance, at least about 0.5 % by weight of the protein, and at least about 25 mOsm for an osmolarity of the drying-salt solution.

Embodiments of the present invention additionally provide a method for reconstituting dried cells comprising drying solute-loaded cells in a drying solution having a salt solution (e.g., PBS), a solute (e.g., trehalose), an inert substance (e.g., a starch), and a protein (e.g., albumin) to produce dried cells, and reconstituting the dried cells in a rehydration solution having the salt solution, the solute, the inert substance, and the protein to produce reconstituted cells. The drying solution may comprise at least about 50 mM of the solute, at least about 25 mOsm osmolarity of the salt solution,

at least about 2.0 % by weight of the inert substance, and at least about 0.5 % by weight of the protein. The rehydration solution may comprise at least about 50 mM of the solute, at least about 25 mOsm osmolarity of the salt solution, at least about 2.0 % by weight of the inert substance, and at least about 0.5 % by weight of the protein. The dried cells may comprise from about 25 mM to about 300 mM of the solute, from about 5 mOsm to about 100 mOsm osmolarity for the salt solution, from about 0.1 % by weight to about 2.5 % by weight of the protein, and from about 1.0 % by weight to about 15.0 % by weight of the inert substance. The dried cells may comprise from about 60 mM to about 80 mM trehalose, from about 10 mOsm to about 40 mOsm PBS, from about 0.3 % by weight to about 9.0 % by weight albumin, and about 1.0 % by weight to about 4.0 % by weight starch.

These provisions, together with the various ancillary provisions and features which will become apparent to those skilled in the art as the following description proceeds, are attained by the processes and cells of the present invention, preferred embodiments thereof being shown with reference to the accompanying drawings, by way of example only, wherein:

Brief Description of the Drawings

In the drawings:

Figure 1 graphically illustrates the loading efficiency of trehalose plotted versus incubation temperature of human platelets.

Figure 2 graphically illustrates the loading efficiency (cytosolic concentration divided by the extracellular concentration, the sum multiplied by 100) following incubation as a function of incubation time;

Figure 3 graphically illustrates the internal trehalose concentration of human platelets versus external trehalose concentration as a function of temperature at a constant incubation or loading time.

Figure 4 graphically illustrates the loading efficiency of trehalose into human platelets as a function of external trehalose concentration.

Figure 5 graphically illustrates intracellular trehalose concentration of human erythrocytes as a function of extracellular trehalose at respective temperatures of 4° C and 37° C.

Figure 6 graphically illustrates the fragility index of erythrocytic cells incubated overnight at respective temperatures of 4° C and 37° C in the presence of and as a function of increasing intracellular trehalose concentrations.

Figure 7 graphically illustrates trehalose uptake (i.e., intracellular trehalose mM) and hemolysis (i.e., % hemolysis) as a function of incubation temperature (°C).

Figure 8 graphically illustrates trehalose uptake (i.e., intracellular trehalose mM) as a function of the osmolarity of the washing buffer.

Figure 9 graphically illustrates % hemolysis of loaded cells vs. time (hours) of incubation of the loaded 300 mOsm PBS buffer.

Figure 10 graphically illustrates time course (incubation time, hours) of hemolysis(%) of trehalose loaded cells as a function of the trehalose concentration in the incubation buffer.

Figure 11 graphically illustrates time course (incubation time, hours) of hemolysis(%) of trehalose loaded cells as a function of the composition of the incubation buffer, and illustrating that HES and albumin (HSA) do not have any detrimental effect on cell hemolysis during incubation.

Figure 12 is a 40X picture of rehydrated erythrocytic cells having no intracellular trehalose prior to freeze-drying.

Figure 13 is a 40X picture of rehydrated erythrocytic cells having 3 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA.

Figure 14 is a 40X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA.

Figure 15 is a 100X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA.

Figure 16 is a graph of hemolysis(%) of trehalose loaded, freeze-dried and rehydrated erythrocytic cells as a function of intracellular trehalose concentration (mM), graphically illustrating the effect of cytoplasmic trehalose on the survival of rehydrated erythrocytic cells.

Figure 17 is a graph of mean corpuscular hemoglobin of trehalose-loaded, freeze-dried and rehydrated erythrocytic cells as a function of intracellular trehalose concentration (mM), graphically illustrating that as the concentration of intracellular trehalose increases for rehydrated erythrocytic cells, the mean corpuscular hemoglobin (the amount of hemoglobin found in intact erythrocytic cells) also increases for rehydrated erythrocytic cells.

Figure 18 is a graph showing the ATP level of erythrocytes in buffers with different compositions during 5 hours incubation at 38-41°C.

Figure 19 is a graph showing the level of 2,3-DPG during 5 hours incubation at 38-41°C in buffers with different composition.

Figure 20 is a graph showing the effect of pre-hydration time on the survival of freeze-dried and rehydrated erythrocytes.

Figure 21 is a graph illustrating the results of having studied the effects of α -crystallin on the percent hemolysis.

Figure 22 is a graph illustrating the combined effect of α -crystallin, Zn^{2+} ions and pre-hydration on the survival of erythrocytic cells.

Figure 23 is a graph illustrating the effect of rejuvenating buffer on the synthesis of ATP and 2,3-DPG in rehydrated erythrocytes.

Detailed Description of Preferred Embodiments of the Invention

Compositions and embodiments of the invention include methods for loading solutes into cells, as well as cells that have been manipulated (e.g., by freeze-drying) or modified (e.g., loaded with a chemical or drug) in accordance with methods of the present invention. The cells may be any type of cell including, not by way of limitation, erythrocytic cells, eukaryotic cells or any other cell, whether nucleated or non-nucleated.

The term "erythrocytic cell" is used to mean any red blood cell. Mammalian, particularly human, erythrocytes are preferred. Suitable mammalian species for providing erythrocytic cells include by way of example only, not only human, but also equine, canine, feline, or endangered species.

The term "eukaryotic cell" is used to mean any nucleated cell, i.e., a cell that possesses a nucleus surrounded by a nuclear membrane, as well as any cell that is derived by terminal differentiation from a nucleated cell, even though the derived cell is not nucleated. Examples of the latter are terminally differentiated human red blood cells. Mammalian, and particularly human, eukaryotes are preferred. Suitable mammalian species include by way of example only, not only human, but also equine, canine, feline, or endangered species.

Molarity, or millimolarity, mM, is the number of moles (or millimoles) of a solute per liter of solution and is a measure of the concentration. Osmolarity (Osm), or milliosmolarity (mOsm), is a count of the number of dissolved particles per liter of solution and is a measure of the osmotic pressure exerted by solutes. Biological membranes, such as cell membranes, can be semi-permeable because they allow water and some small molecules to pass, but block the passage of proteins or macromolecules. Since the osmolarity of a solution is equal to the molarity times the number of particles per molecule, 600 mM trehalose is equal to 600 mOsm trehalose because trehalose does not dissociate in water. However, with respect to compounds that dissociate in water, such as NaCl, 1 mM NaCl is equal to 2 mOsm NaCl because it has two particles. Similarly, 100 mM NaCl is equal to 200 mOsm NaCl. Thus, for a 300 mOsm PBS buffer (100 mM NaCl, 9.4 mM Na_2HPO_4 , 0.6 mM KH_2PO_4 , pH 7.4), 300 mOsm refers to all of the osmotically active particles in the PBS solution, with 200 mOsm of the 300 mOsm stemming from NaCl.

Broadly, the preparation of solute-loaded cells in accordance with embodiments of the invention comprises the steps of loading one or more cells with a solute by placing one or more cells in a solute solution having a solute concentration of sufficient magnitude to produce hyperosmotic pressure on the cell for transferring the solute from the solution into the cell. For increasing the transfer or uptake of the solute from the solute solution, the solute solution temperature or incubation temperature has a temperature above about 25°C, more preferably above 30°C, such as from about 30°C to about 40°C. In another embodiment of the invention, a solute solution (e.g., trehalose solution) has a solute (e.g., trehalose) concentration of at least about 25 %, preferably at least about 50 %, greater

than the intracellular osmolarity of the cells for loading the solute into the cells.

For various embodiments of the invention, a solute solution has a solute concentration ranging from about 25 % to at least about 1000 % greater than the intracellular osmolarity of the cell. For additional various embodiments of the invention, the solute solution, especially when the solute solution is employed as a loading buffer, has a solute concentration ranging from about 320 mM to about 4000 mM, preferably from about 320 mM to about 2000 mM, more preferably from about 500 mM to about 1000 mM.

The method(s) for various embodiments of the present invention may additionally comprise preventing a decrease in a loading gradient and/or a loading efficiency gradient in the loading of the solute into the cells. Preventing a decrease in a loading efficiency gradient in the loading of the solute into the cells comprises maintaining a positive gradient of loading efficiency (e.g., in %) to concentration (e.g., in mM) of the solute in the solute solution. Preventing a decrease in a loading gradient in the loading of the solute into the cells comprises maintaining a concentration of the solute in the solute solution below a certain concentration (e.g., below a concentration ranging from about 35 mM to about 65 mM, more particularly below from about 40 mM to about 60 mM, or below from about 45 mM to about 55 mM, such as below about 50 mM); and/or maintaining a positive gradient of concentration of solute loaded into the cells to concentration of the solute in the solute solution.

The solute solution for various embodiments of the present invention may be used for loading and/or washing and/or freeze-drying and/or rehydration, or for any other suitable purpose. When the solute solution is employed for loading a solute into

the cells, the solute solution may be any suitable physiologically acceptable solution in an amount and under conditions effective to cause uptake or "introduction" of the solute from the solute solution into the cells. A physiologically acceptable solution is a suitable solute-loading buffer, such as any of the buffers stated in the previously mentioned related patent applications, all having been incorporated herein by reference thereto. The solute solution may also be any suitable physiologically acceptable solution in an amount and under conditions effective for washing and/or freeze drying and/or rehydration. Therefore, the solute solution may be used as a washing buffer for washing loaded cells and/or as a freeze-drying buffer for freeze-drying loaded cells and/or as a rehydration buffer for rehydrating thawed cells in reconstituting cells. Thus, any of the solute solutions for embodiments of the present invention may be used for any suitable purpose, including loading, washing, freeze-drying, and rehydration.

For particular embodiments of the present invention, especially when the solute solution is being employed as a loading buffer and/or washing buffer, the solute solution comprises a solute and a salt solution. In other particular embodiments of the invention, especially when the solute solution is being employed as a freeze-drying buffer and/or a rehydration buffer, the solute solution comprises a salt solution (e.g., PBS), a protein, a solute, and at least one inert substance. However, it is to be understood that the solute solution comprising a salt solution, a protein, a solute, and an inert substance may be used for any other suitable purpose including for loading a solute into cells and for washing solute-loaded cells.

Protein, when referred to herein, means any suitable protein (e.g., simple or conjugated protein), including any complex, high polymer containing carbon, hydrogen, oxygen, nitrogen, and usually sulfur, and composed of chains of amino acids connected by peptide linkages. Protein includes albumin, which when referred to herein means any suitable albumin (e.g., bovine albumin), including any of a group of water-soluble proteins of wide occurrence in such natural products as milk (lactalbumin), blood serum, eggs (ovalbumin). Preferably, the albumin comprises human serum albumin (HSA).

The solute is preferably a carbohydrate (e.g., an oligosaccharide) selected from the following groups of carbohydrates: a monosaccharide (e.g., bioses, trioses, tetroses, pentoses, hexoses, heptoses, etc), a disaccharide (e.g., lactose, maltose, sucrose, melibiose, trehalose, etc), a trisaccharide (e.g., raffinose, melezitose, etc), or tetrasaccharides (e.g., lupeose, stachyose, etc), and a polysaccharide (e.g., dextrans, starch groups, cellulose groups, etc). More preferably, the solute is a disaccharide, with trehalose being the preferred, particularly since it has been discovered that trehalose does not degrade or reduce in complexity upon being loaded. Thus, in the practice of various embodiments of the invention, trehalose is transferred from a solution into the cells without degradation of the trehalose.

The salt solution may be any suitable physiologically acceptable solution in an amount and under conditions effective to function as a carrier medium for a solvent, or for a mixture of a solvent, a protein and/or an inert substance. The salt solution may comprise a phosphate buffered saline (PBS) solution comprising NaCl, Na_2HPO_4 , and KH_2PO_4 . A suitable PBS buffer is 100 mOsm PBS buffer (51.3 mM NaCl, 1.87 mM Na_2HPO_4 , 0.35 mM KH_2PO_4 , pH 7.2).

The inert substance is preferably a carbohydrate, such as any of the carbohydrates previously mentioned above.

Preferably, the inert substance comprises a polysaccharide.

More preferably, the inert substance comprises a starch, such as, by way of example only, hydroxy ethyl starch (HES).

The quantities of solute, protein and inert substance employed in the solute solution, more specifically in combination with a saline solution, are of suitable quantities and proportion for minimizing the loss or destruction of cells, more particularly for minimizing hemolysis, especially after freeze-drying and reconstitution (e.g., prehydration and rehydration), and/or especially when the solute solution is employed as a freeze-drying buffer and/or rehydration buffer.

For various embodiments of the present invention, the solute solution comprises: a solute and a salt solution. The concentration of the solute in the solute solution may be at least about 50 mM, such as ranging from about 50 mM to about 3000 mM, preferably from about 100 mM to about 1500 mM, more preferably from about 150 mM to about 1000 mM, most preferably from about 200 mM to about 600 mM. The osmolarity of the salt solution may be at least about 25 mOsm, such as ranging from about 25 mOsm to about 1000 mOsm, preferably from about 50 mOsm to about 300 mOsm, more preferably from about 75 mOsm to about 200 mOsm. The solute solution comprising a solute and a salt solution may be used for any suitable purpose including as a loading buffer and/or as a washing buffer.

For additional various embodiments of the present invention, the solute solution may further comprise (in addition to the solute and the salt solution) a protein and/or an inert substance. The amount or quantity of the inert substance (e.g., HES) in the solute solution may be at least about 2.0 % by

weight, such as ranging from about 2.0 % by weight to about 50 % by weight, preferably from about 5 % by weight to about 35 % by weight, more preferably from about 10 % by weight to about 30 % by weight, most preferably from about 12 % by weight to about 20 % by weight (e.g., about 15 % by weight). The amount or quantity of the protein (e.g. HSA) in the solute solution may be at least about 0.5 % by weight, such as ranging from about 0.5 % by weight to about 15 % by weight, preferably from about 1 % by weight to about 10 % by weight, more preferably from about 1.5 % by weight to about 8 % by weight, most preferably from about 1.5 % by weight to about 5 % by weight (e.g., about 2.5 % by weight). The solute solution comprising a solute, a salt solution, a protein and/or an inert substance may be used for any suitable purpose including as a freeze drying buffer and/or rehydration buffer.

An extracellular medium of about 280-320 mOsm is considered iso-osmotic for cells, particularly erythrocytic cells, with regard to the amount of permeable solutes in the cytoplasm. Any increase of the amount of solutes in the extracellular medium creates an osmotic shock, ranging from a mild shock at about 350 mM trehalose to a strong shock at about 420 mM trehalose, and a leakage of water which would reversibly reduce the cell volume. However, small molecular weight solutes, such as trehalose, in an extracellular medium in a concentration higher than about 320 mM, can pass through the membrane of a cell using a diffusion vector. It has been discovered that an extracellular concentration of trehalose higher than about 450 mM (or mOsm), which is about 50% greater than an intracellular milliosmolarity, will produce an osmotic shock that will result in trehalose uptake. Increasing the extracellular trehalose

concentration leads to even higher osmotic shock and higher trehalose uptake.

Other embodiments of the present invention provide for retaining a solute in a cell. Preferably, after the cells have been loaded with a solute, such as an oligosaccharide (e.g., trehalose), the cells are then washed. More preferably, during the washing of the cells the solute is retained in the cells. Washing leads to hemolysis of the fragile cells and removal of cellular fragments and free hemoglobin. The net result is that the remaining cells do indeed have an elevated trehalose content. The washing may be with a washing solution (e.g., such as a washing buffer having an oligosaccharide), and retention of the solute in the cell increases from about 25% to about 175% when a buffer concentration (e.g., the osmolarity of all osmotically active particles within the washing buffer solution) increases from about 50% to about 400%, more preferably from about 50% to about 150% when a buffer concentration increases from about 100% to about 300%, and most preferably from about 75% to about 125% (e.g., about 100%) when a buffer concentration increases from about 150% to about 250% (e.g., about 200%). The washing of the cell with a washing buffer includes employing a ratio of a buffer concentration (mOsm) (e.g., an extracellular buffer concentration) to an intracellular solute concentration (mM) ranging from about 14.0 to about 4.0, such as from about 12.0 to about 5.0, including from about 9.0 to about 6.0 and from about 8.0 to about 7.0 (e.g., about 7.5).

As indicated in patent application Serial No. 10/052,162, which claims the benefit of patent application Serial No. 09/501,773, filed February 10, 2000, with respect to common subject matter, the amount of the preferred trehalose loaded inside the cells ranges from about 10 mM to about 50 mM, and is achieved by incubating the cells to preserve biological

properties during freeze-drying with a trehalose solution, preferably a trehalose solution that has up to about 50 mM trehalose therein. Higher concentrations of trehalose during incubation are not preferred, particularly since an embodiment of the invention includes preventing a decrease in a loading gradient, or a loading efficiency gradient, in the loading of the solute into the cell. It has been discovered that preventing a decrease in a loading gradient, or a loading efficiency gradient, in the loading of a solute (e.g., an oligosaccharide, such as trehalose) into a cell comprises maintaining a concentration of the solute in the solute solution below a certain concentration (e.g., below about 75 mM, such as below about a concentration ranging from about 35 mM to about 65 mM, more particularly below from about 40 mM to about 60 mM, or below from about 45 mM to about 55 mM, such as below about 50 mM). It has been further discovered that preventing a decrease in a loading gradient, or a loading efficiency gradient, in the loading of a solute into a cell comprises maintaining a positive gradient of loading efficiency to concentration of the solute in the solute solution.

As further indicated in co-pending patent application Serial No. 10/052,162, the effective loading of trehalose is also accomplished by means of using an elevated temperature of from greater than about 25° C to less than about 40° C, more preferably from about 30°C to less than about 40°C, most preferably about 37°C. This is due to the discovery of the second phase transition for cells.

Referring now to Fig. 1, there is seen a graphical illustration from co-pending patent application Serial No. 10/052,162 of the loading efficiency of trehalose plotted versus incubation temperature of human platelets. The trehalose loading

efficiency begins a steep slope increase at incubation temperatures above about 25°C and continues up to about 40°C. The trehalose concentration in the exterior solution (that is, the solute solution or loading buffer) and the temperature during incubation together lead to a trehalose uptake that occurs through fluid phase endocytosis. Example 1 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 1. It is believed that the graphical illustration of the loading efficiency in Fig. 1 could be generally applicable for cells in general.

Referring now to Fig. 2, there is seen an illustration from co-pending patent application Serial No. 10/052,162 of trehalose loading efficiency for human blood platelets as a function of incubation time. More specifically, Fig. 2 is a graphical illustration of the loading efficiency (cytosolic concentration divided by the extracellular concentration, the sum multiplied by 100) following incubation as a function of incubation time. Example 1 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 2. It is believed that the graphical illustration of the loading efficiency in Fig. 2 could also be generally applicable for cells in general.

Referring now to Figure 3, there is seen a graphical illustration from patent application Serial No. 10/052,162 of the internal trehalose concentration of human platelets versus external trehalose concentration as a function of 4° C and 37° C temperatures at a constant incubation or loading time. In Figure 4 there is seen a graphical illustration from patent application Serial No. 10/052,162 of the loading efficiency of trehalose into human platelets as a function of external trehalose

concentration. Example 1 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figures 3 and 4. In additional embodiments of the present invention, it is further believed that the general findings illustrated in Figs. 3 and 4 with respect to platelets are generally broadly applicable to cells in general.

Thus, applying the findings illustrated in Fig. 3 and in Fig. 4 to solutes and cells in general, a decrease in a loading gradient or a loading efficiency gradient in the loading of a solute into a cell may be prevented. For an embodiment of the present invention and as broadly illustrated in Figure 3, preventing a decrease in a loading gradient or a loading efficiency gradient in the loading of the solute (e.g., an oligosaccharide such as trehalose) into the cell comprises maintaining a concentration of the solute (e.g., an oligosaccharide such as trehalose) in the solute solution (e.g. an oligosaccharide solution such as a trehalose solution) below a solute concentration ranging from about 35 mM to about 65 mM, more specifically a solute concentration ranging from about 40 mM to about 60 mM, more specifically further a solute concentration ranging from about 45 mM to about 55 mM (e.g., about 50 mM). In another embodiment of the present invention and as best illustrated in Figure 4, preventing a decrease in a loading gradient or a loading efficiency gradient in the loading of the solute (e.g., an oligosaccharide, such as trehalose) into the cell comprises maintaining a positive gradient of loading efficiency (e.g., loading efficiency in %) to concentration (e.g., concentration in mM) of the solute in the solute solution (e.g. an oligosaccharide solution, such as a trehalose solution).

When a solute is loaded from a solute solution into one or more cells, the solute solution preferably has a solute concentration of sufficient magnitude to produce hyperosmotic pressure on the one or more cells. It has been discovered that the basis for the loading of the solute into the cells is dependent upon osmotic shock. The magnitude of osmotic shock and hyperosmotic pressure on the cells depends on the difference between internal solute concentration, or the intracellular osmolarity, within the cells, and the external solute concentration within the solute solution, or the extracellular cellular solute concentration. For embodiments of the invention, the solute solution has a solute concentration ranging from about 320 mM to about 4000 mM, preferably from about 320 mM to about 2000 mM, more preferably from about 500 mM to about 1000 mM.

In another embodiment of the present invention, the solute solution, especially when used for loading the solute into one or more cells, may comprise a solute and a salt solution having a suitable osmolarity (mOsm). Preferably, the ratio of the osmolarity (mOsm) of the salt solution to the solute concentration (mM) in the solution ranges from about 0.04 to about 1.0, preferably from about 0.05 to about 0.50, more preferably from about 0.07 to about 0.30, most preferably from about 0.10 to about 0.20. The osmolarity (mOsm) of the solute solution for these embodiments of the invention is the osmolarity of the osmotically active particles except, or other than, the osmolarity of the solute. As indicated previously, the osmolarity of the solute solution may range from about 25 mOsm to about 1000 mOsm, preferably from about 50 mOsm to about 300 mOsm, more preferably from about 75 mOsm to about 200 mOsm. As indicated previously, the solute solution may be any suitable

solution for purposes for embodiments of the present invention. Preferably, the solute solution comprises a salt solution, such as a phosphate buffered saline (PBS) comprising NaCl, Na_2HPO_4 , and KH_2PO_4 . A suitable PBS buffer is 100 mOsm PBS buffer (51.3 mM NaCl, 1.87 mM Na_2HPO_4 , 0.35 mM KH_2PO_4 , pH 7.2).

It has also been discovered that the basis for the loading of the solute into the cells is not only dependent upon osmotic shock, but is also dependent upon the thermal effects on flux of the solute across the membranes of the cells. The higher the thermal effects on flux of the solute across the membranes of the cells, the larger the amount of solute loaded into the cells. Stated alternatively, loading of a solute into cells increases as the temperature of the solute solution increases. Referring now to Figure 5, there is seen a graphical illustration of intracellular trehalose concentration as a function of extracellular trehalose at respective temperatures of 4°C and 37°C . Thus, at a temperature ranging from about 30°C to about 40°C (e.g. at about 37°C) a gradient of a solute concentration (M), such as an oligosaccharide (e.g., trehalose) concentration, within a cell (e.g., an erythrocytic cell) to extracellular solute concentration (M) within a loading solution (or buffer) ranges from about 0.130 to about 0.200. At a temperature ranging from about 0°C to about 10°C (e.g. at about 4°C) a gradient of a solute concentration (M), such as an oligosaccharide (e.g., trehalose) concentration, within a cell (e.g., an erythrocytic cell) to extracellular solute concentration (M) within a loading solution (or buffer) ranges from about 0.04 to about 0.12, more specifically from about 0.04 to about 0.08, and from about 0.08 to about 0.12, depending on the quantity of extracellular solute concentration. Example 2 below provides the more specific testing conditions and

parameters which produced the graphical illustrations of Figure 5.

Referring now to Figure 6, there is seen a graphical illustration of the fragility index of erythrocytic cells incubated overnight at respective temperatures of 4° C and 37° C in the presence of and as a function of increasing intracellular trehalose concentrations. The osmotic fragility index was generated by the extent of hemolysis as a function of the NaCl concentration. The graphical illustration of Figure 6 represents a test for investigating the effects of hyperosmotic treatment rendering erythrocytic cells more sensitive to change in intracellular osmolarity. NaCl was loaded into erythrocytic cells from a 100 mOsm PBS buffer at loading 100 mOsm PBS buffer temperatures of 4° C and 37° C for extracellular trehalose concentrations of 0 mM (control cells), 250 mM, 500 mM, 600 mM, 700 mM, 800 mM and 1000 mM. Data blocks, respectively generally indicated as **60** and **62**, represent the intracellular trehalose concentrations for 100 mOsm PBS solution loading temperatures of 4° C and 37° C. The mOsm/kg values of NaCl represent extracellular NaCl osmolarity of the erythrocytic cells resulting from the transfer of NaCl from the PBS loading buffer into the erythrocytic cells. The erythrocytic cells that had been loaded in trehalose solutions (between 250 mM and 1000 mM) in 100 mOsm PBS were suspended in increasing concentrations of NaCl (between 50 and 600 mOsm NaCl). The percent hemolysis measured after resuspending the loaded cells in NaCl represents the fragility index. The data show that the erythrocytic cells were stable osmotically in trehalose media with concentrations between 250 mM and 800 mM trehalose at both 37° C and 4° C. In 1000 mM trehalose at 37° C, there is a high increase in the fragility index suggesting that the cells were unstable in this

medium (1000mM trehalose in 100 mOsm PBS). Clearly, at moderate intracellular concentrations of trehalose, osmotic fragility as measured by a standard assay was not severely altered. Thus, erythrocytic cells may be loaded with trehalose concentrations up to about 900 mM (i.e., a trehalose concentration between 800 mM and 1000 mM). Example 3 below provides specific testing conditions and parameters which produced the graphical illustrations of Figure 6.

Thus, from the findings graphically illustrated in Figs. 5 and 6, and as more fully explained in Examples 2 and 3 below, temperature of a solute loading solution has an effect in loading a solute from a solute solution into a cell. The effects of temperature, as well as cellular hemolysis, of a trehalose loading solution in loading of trehalose into a cell was tested. The test results are illustrated in Figure 7, which is a graphical illustration of trehalose uptake (i.e., intracellular trehalose mM) and hemolysis (i.e., % hemolysis) as a function of incubation temperature ($^{\circ}\text{C}$). The incubation time was about 6 hours and the medium contained about 800 mM trehalose/100 mM PBS. Figure 7 illustrates that effective loading occurs above 30°C , and that as the loading temperature of the trehalose loading solution increases, there is slight hemolysis. Example 4 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 7.

As previously indicated, after a cell (e.g., an erythrocytic cell) has been loaded with a solute (e.g., trehalose), further embodiments of the present invention provide for retaining the solute in the cells. One means for retaining solute within solute-loaded cells is to wash the cells, more specifically by washing the cells and retaining the solute in the cells during the washing. As also previously indicated, the

washing of the cells is preferably with a washing buffer. It has been discovered that retention of the solute in the cells increases from about 25% to about 175% when a buffer concentration (e.g., the osmolarity of all osmotically active particles within the washing buffer solution) increases from about 50% to about 400%, more preferably from about 50% to about 150% when a buffer concentration increases from about 100% to about 300%, and most preferably from about 75% to about 125% (e.g., about 100%) when a buffer concentration increases from about 150% to about 250% (e.g., about 200%). It has been further discovered that the washing of the cells with a washing buffer includes employing a ratio of an extracellular buffer concentration (mOsm) to an intracellular trehalose concentration (mM) ranging from about 14.0 to about 4.0, more particularly from about 12.0 to about 5.0, including from about 9.0 to about 6.0 and from about 8.0 to about 7.0 (e.g., about 7.5). Thus, because solute loaded cells are hyperosmotic to a washing buffer, increasing the extracellular osmolarity increases retention of the solute, particularly during washing of the cells, as shown in Figure 8 which graphically illustrates intracellular trehalose concentration (mM) as a function of the osmolarity of the washing buffer. As shown in Figure 8, when the extracellular buffer concentration was increased from 300 mOsm PBS to 900 mOsm PBS during washing, the final intracellular trehalose concentration doubled. The 300 mOsm PBS had no trehalose concentration, and the 900 mOsm PBS also had no trehalose concentration. Example 5 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 8.

After the cells have been effectively loaded with a solute and subsequently washed, the cells may then be contacted with a drying buffer. The drying buffer should include the solute,

preferably in amounts up to about 100 mM. The solute in the drying buffer assists in spatially separating the cells as well as stabilizing the cell membranes on the exterior. The drying buffer preferably also includes a bulking agent (to further separate the cells). Albumin may serve as a bulking agent, but other polymers may be used with the same effect. If albumin is used, it is preferably from the same species as the cells. Suitable other polymers, for example, are water-soluble polymers such as HES (hydroxy ethyl starch) and dextran.

For other embodiments of the present invention, and as previously mentioned, the solute solution may serve as the drying buffer. The solute solution, when functioning as a drying buffer, may comprise at least about 50 mM of the solute, at least about 2.0 % by weight of the inert substance, at least about 0.5 % by weight of the protein, and at least about 25 mOsm for an osmolarity of the salt solution. More specifically, as indicated, the solute solution for drying buffer purposes may comprise the solute having a concentration ranging from about 50 mM to about 3000 mM, preferably from about 100 mM to about 1500 mM, more preferably from about 150 mM to about 1000 mM, most preferably from about 200 mM to about 600 mM. The osmolarity of the salt solution in the solute solution may range from about 25 mOsm to about 1000 mOsm, preferably from about 50 mOsm to about 300 mOsm, more preferably from about 75 mOsm to about 200 mOsm. The amount or quantity of the inert substance (e.g., HES) in the solute solution may range from about 2.0 % by weight to about 50 % by weight, preferably from about 5 % by weight to about 35 % by weight, more preferably from about 10 % by weight to about 30 % by weight, most preferably from about 12 % by weight to about 20 % by weight (e.g., about 15 % by weight). The amount or quantity of the protein (e.g. HSA) in the

solute solution may range from about 0.5 % by weight to about 15 % by weight, preferably from about 1 % by weight to about 10 % by weight, more preferably from about 1.5 % by weight to about 8 % by weight, most preferably from about 1.5 % by weight to about 5 % by weight (e.g., about 2.5 % by weight).

The solute loaded cells in the drying buffer may then be dried while simultaneously cooled to a temperature below about -32°C . A cooling, that is, freezing, rate is preferably between -30°C and $-1^{\circ}\text{C}/\text{min}$. and more preferably between about $-2^{\circ}\text{C}/\text{min}$ to $-5^{\circ}\text{C}/\text{min}$. Drying may be continued until about 95 weight percent of water has been removed from the cells. During the initial stages of lyophilization, the pressure is preferably at about $10 \times$

10^{-6} torr. As the samples dry, the temperature can be raised to be warmer than -32°C . Based upon the bulk of the sample, the temperature and the pressure it can be empirically determined what the most efficient temperature values should be in order to maximize the evaporative water loss. Freeze-dried cell compositions preferably have less than about 5 weight percent water.

After freeze drying and storage of the cells, the process of using such a dehydrated cell composition comprises rehydrating the cells. The rehydration preferably includes a prehydration step, sufficient to bring the water content of the freeze-dried cells to between about 20 weight percent and about 50 percent, preferably from about 20 weight percent to about 40 weight percent. More preferably, when reconstitution of the freeze dried cells is desired, the freeze dried cells are prehydrated in moisture saturated air at about 37°C for about one hour to about three hours, followed by rehydration. Use of prehydration yields cells with a much more dense appearance and

with no balloon cells being present. The preferred prehydration step brings the water content of the freeze-dried cells to between about 20 weight percent to about 50 weight percent. Rehydration of the prehydrated cells may be with any aqueous based solutions, depending upon the intended application.

Referring now to Figure 9, there is seen a graphical illustration of hemolysis (%) of loaded cells as a function of the time (in hours) in 300 mOsm PBS. The cells were loaded with 700 mM trehalose in 100 mM PBS. As illustrated in Figure 9, the % hemolysis of loaded cells is below about 10 % (i.e., about 7 %) if the loaded cells are stored in the 300 mOsm PBS within or less than about 3 hours (e.g., within or less than about 2 hours) after being loaded, preferably after being loaded and subsequently washed. The % hemolysis of the loaded cells precipitously increases (i.e., greater than about 10 % hemolysis) if more than 3 hours lapses after loading and/or washing loaded cells and incubating them in 300 mOsm PBS. Example 6 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 9.

Hemolysis of the cells not only depends on the period of time of subsequently freeze-drying after loading the cells, or after washing loaded cells, but also on the incubation time and the quantity of intracellular solute loaded in the cells, as well as, in some cases, on whether or not the inert substance and/or protein has been admixed with the solute in the salt solution, and/or, in other cases, the quantity of the inert substance and/or the protein that has been added. Referring now to Figure 10, there is seen a graphical illustration of time course (incubation time, hours) of hemolysis(%) of trehalose-loaded cells as a function of the composition of the incubation

buffer. Percent (%) hemolysis represents the hemolysis following incubation in a designated incubation buffer, which typically, may be different from a loading buffer. Figure 10 represents an illustration of the % hemolysis of trehalose-loaded cells in buffers with different concentration of trehalose. Broadly, as the concentration of the solute (e.g., trehalose) increases, hemolysis decreases. Curve **1004** represents % hemolysis vs. incubation time (hours) when incubating cells with no trehalose and a 300 mM PBS salt solution. Curve **1008** represents % hemolysis vs. incubation time (hours) when incubating cells with 100 mM_trehalose and a 300 mM PBS salt solution. Curve **1012** represents % hemolysis vs. incubation time (hours) when incubating cells with 200 mM trehalose and a 300 mM PBS salt solution. Curve **1016** represents % hemolysis vs. incubation time (hours) when incubating cells with 300 mM_trehalose and a 300 mM PBS salt solution. Thus, broadly, when trehalose-loaded cells are incubated in an incubation period (e.g. from 0 hours to about 3 hours) with a solution comprising a salt solution (e.g., 300 mM PBS) and at least about 200 mM of a solute (e.g., trehalose), hemolysis of the cells is reduced to less than about 10 %, more preferably to less than about 5%. Example 7 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 10.

Referring now to Figure 11, there is seen a graphical illustration of the time course (incubation time, hours) of hemolysis(%) of trehalose-loaded cells as a function of the composition of the incubation buffer, and illustrating that HES and albumin (HSA) do not have any detrimental effect on cell hemolysis during incubation. Figure 11 illustrates % hemolysis of trehalose-loaded cells in buffers with different amount of HES and HSA for determining which buffers provide highest cell

stability, assessed as the lowest % hemolysis. As indicated, hemolysis of the cells not only depends on the period of time of subsequently freeze-drying after loading the cells, or after washing loaded cells, but also, in some cases, on whether or not the inert substance and/or protein has been admixed with the solute in the salt solution, and/or, in other cases, the quantity of the inert substance and/or the protein that has been added. Curve **1110** represents hemolysis vs. incubation time (hours) when incubating cells with 300 mOsm PBS. Curve **1114** represents hemolysis vs. incubation time (hours) when incubating cells with 15 % HES. Curve **1118** represents hemolysis vs. incubation time (hours) when incubating cells with 30 % HES. Curve **1122** represents hemolysis vs. incubation time (hours) when incubating cells with 2.5% HSA. Curve **1126** represents hemolysis vs. incubation time (hours) when incubating cells with 5% HSA. Curve **1130** represents hemolysis vs. incubation time (hours) when incubating cells with 2.5% HAS and 15% HES. Thus, adding the inert substance and/or the protein, particularly in the quantities of 15% (i.e, HES) and 2.5 % (i.e., HSA), to the incubation solution assists in further reducing hemolysis. As will be further explained below, the addition of the inert substance and/or the protein, both in the freeze-drying buffer and in the rehydration buffer, has significantly improved cell drying (e.g., freeze-drying) and rehydration such that rehydrated cells have normal discoid morphology. Example 8 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 11.

Referring now to Figures 12-15, there are seen pictures of cells (e.g., erythrocytic cells) which illustrate that intracellular trehalose, more particularly intracellular trehalose along with intracellular inert substance and protein,

improve the survival of rehydrated cells. Figure 12 is a 40X picture of rehydrated erythrocytic cells having no intracellular trehalose prior to freeze-drying. Figure 13 is a 40X picture of rehydrated erythrocytic cells having 3 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Figure 14 is a 40X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Figure 15 is a 100X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Example 9 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figures 12-15.

The concentration of intracellular trehalose in rehydrated cells is also important for subsequent stabilization of the rehydrated cells. Referring now to Figure 17, there is seen a graph of mean corpuscular hemoglobin of trehalose-loaded, freeze-dried and rehydrated erythrocytic cells as a function of intracellular trehalose concentration (mM). Figure 17 illustrates that as the concentration of intracellular trehalose increases for rehydrated erythrocytic cells, the mean corpuscular hemoglobin (MCH, the amount of hemoglobin found in intact erythrocytic cells) also increases for rehydrated erythrocytic cells. Example 11 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 17. An intracellular trehalose concentration (mM) of about 40 mM and (or to) about 42 mM produces an MCH of about 9 (pg). When the intracellular

trehalose concentration (mM) increases to about 60 mM, the MCH precipitously increases to above about 10 (pg), more specifically above or greater than about 14 (pg). Thus, embodiments of the present invention include loading cells with an effective amount of solute for stabilizing cells. An effective amount of a solute is greater than about 50 mM, such as 60 mM or above. Furthermore, as previously indicated, as intracellular concentration increases, there is a significant decrease in the percent (%) hemolysis, to an extent of less than about 10 %, and even less than about 5 %, as broadly illustrated in Figure 16 which is a graph of hemolysis(%) of trehalose loaded erythrocytic cells as a function of intracellular trehalose concentration (mM), graphically illustrating the effect of cytoplasmic trehalose on the survival of rehydrated erythrocytic cells. Example 10 below provides the more specific testing conditions and parameters which produced the graphical illustrations of Figure 16. As cytoplasmic intracellular trehalose approaches a concentration of about 100 mM, % hemolysis falls to below about 10 %.

The following protocol has been discovered as yielding significant survival of freeze-dried cells. The loading buffer comprised about 800 mM trehalose in a salt solution of about 100 mOsm PBS. The incubation time was about 16 hours at a temperature of about 35⁰ C. After the cells were loaded, they were subsequently washed in a washing buffer comprising about 300 mM trehalose in a salt solution of about 100 mOsm PBS. Within about 3 hours after washing the loaded cells, the wash loaded cells were freeze-dried in freeze-drying buffer comprising about 300 mM trehalose, about 100 mOsm PBS, about 2.5 % by wt. HSA, and about 15 % by wt. HES. After freeze-drying, the cells had about 75 mM trehalose, about 25 mOsm PBS, about

0.6 % by wt. HSA and about 4.0 % by wt. HES left in the cells. In various embodiments of the invention for producing maximal survival of the cells, the dried cells comprise from about 25 mM to about 300 mM trehalose, from about 5 mOsm to about 100 mOsm osmolarity for the salt solution, from about 0.1 % by weight to about 2.5 % by weight of the protein, and from about 1.0 % by weight to about 15.0 % by weight of the inert substance; and preferably from about 60 mM to about 80 mM trehalose, from about 10 mOsm to about 40 mOsm PBS, from about 0.3 % by weight to about 9.0 % by weight albumin, and about 1.0 % by weight to about 4.0 % by weight starch. The freeze-dried cells were then reconstituted at about 37° C for about 10 minutes in a rehydration buffer comprising about 188 mM trehalose, about 100 mOsm PBS, about 2.5 % by wt. HSA and about 15.0 % by wt. HES. After rehydration, less than about 5 % of the cells were lysed. Example 12 below also provides the conditions and parameters which produced the foregoing protocol yielding less than 5 % hemolysis.

The following loading protocol has also been discovered as yielding significant survival of freeze-dried cells. The loading protocol includes incubating the erythrocytic cells in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate. ADSOL comprises 111 mM glucose, 2 mM adenine, 154 mM NaCl and 41 mM mannitol. The incubation temperature for loading was between 38 and 41°C, and the time of incubation was 6 hours. This loading procedure yielded lower extent of hemolysis (about 17%), as compared to the hemolysis measured during loading in 800 mM trehalose and 100 mOsm PBS for 16 hours at 37°C. Furthermore, this loading procedure was not accompanied by significant changes in cell morphology. At the same time, the amount of intracellular trehalose was the same as during loading

erythrocytes in 800 mM trehalose and 100 mOsm PBS at 37°C for 16 hours. No washing was applied after termination of the loading step and prior to freeze-drying. Immediately after completing the loading, the cells were mixed gently with the freeze-drying buffer. The final concentration of the freeze-drying buffer was 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% human serum albumin (HSA). The freeze-dried cells were rehydrated at 37°C for about 10 min in a rehydration buffer containing 141 mM trehalose, 75 mOsm PBS 11.25% HES and 1.875% HSA.

During the loading step, the levels of the following two important metabolites were followed as being essential for cell viability: adenosine-3-phosphate (ATP) and 2,3-diphosphoglycerate (2,3-DPG). ATP level correlates with the efficiency of the glycolic pathway which is the major biochemical pathway in erythrocytes. The polyanion 2,3-DPG binds to the central cavity of the hemoglobin tetramer and modulates the affinity of hemoglobin for oxygen. It is important for the oxygen carrying capacity of hemoglobin. The normal level of ATP in freshly isolated erythrocytes was between 3.65 and 4.45 $\mu\text{mole/g Hb}$. Figure 18 shows the ATP level of erythrocytes in buffers with different compositions during 5 hours incubation at 38-41°C. During incubation in 100 mOsm ADSOL and 6.6 mM Na-phosphate (curve **D**) or in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate (curve **B**), the measured ATP level is very similar to that of freshly isolated erythrocytes. When erythrocytes were incubated in 800 mM trehalose and 100 mOsm PBS, the level of ATP was also as high as in fresh cells (curve **E**). It was slightly reduced when cells were incubated in 800 mM trehalose and 100 mOsm ADSOL (without Na-phosphate) (curve **A**), and when the cells were incubated in ADSOL only (462 mOsm) (curve **C**).

Figure 19 presents the level of 2,3-DPG during 5 hours incubation at 38-41°C in buffers with different composition. The normal level of 2,3-DPG in freshly isolated erythrocytes is around 12.8 μ mole/g Hb. The highest 2,3-DPG level was observed in cells incubated in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate (curve B) and in 800 mM trehalose and 100 mOsm ADSOL (curve A). It was decreased for cells incubated in ADSOL (462 mOsm) (curve C), in ADSOL and 6.6 mM Na-phosphate (curve D), in 800 mM trehalose and 100 mOsm PBS (curve E) and in 300 mOsm PBS (curve F).

On the basis of the data in Figures 18 and 19, the incubation medium comprising 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate provides high levels of ATP and 2,3-DPG.

Pre-hydration via exposure to water vapor produces a gradual and more homogenous rehydration of dried biomaterials than direct rehydration. Figure 20 presents the effect of pre-hydration time on the survival of freeze-dried and rehydrated erythrocytes. Erythrocytic cells were loaded in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate at 38-41°C for 6 hours and were freeze-dried in a buffer with a final concentration of 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HSA. Freeze-dried cells were pre-hydrated for various times (between 5 and 30 mins.) and then they were rehydrated at 37°C for 10 min in a buffer containing 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA. The data in Figure 20 show that pre-hydration for 5 min at 37°C provided the lowest percent of hemolysis.

α -crystallin is a member of the small heat shock protein family and is highly abundant in a number of mammalian cell types and tissues. It has been discovered that α -crystallin associates with lipid membranes *in vitro* and preserves their

integrity at high non-lethal temperatures. The results of having studied the effects of α -crystallin on the percent hemolysis are shown in Figure 21. Cells were loaded in either 800 mM trehalose, 100 mOsm ADSOL, 6.6 mM Na-phosphate (Figure 21, data set labeled as 800 mM treh) or in 800 mM trehalose, 100 mOsm ADSOL, 6.6 mM Na-phosphate and 1.2 mg/ml α -crystallin (Figure 21, data set labeled as + α -crystallin 1.2 mg/ml). Cells were subsequently mixed with freeze-drying buffer with final concentration of 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HSA and were freeze-dried. After freeze-drying they were directly rehydrated (no pre-hydration) in 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA. Cells loaded in the presence of 1.2 mg/ml α -crystallin show lower percent hemolysis (49%) in comparison to those loaded without α -crystallin (68%). In the third data set of Figure 21, along with 1.2 mg/ml α -crystallin, there was 0.5 mg/ml α -crystallin added to the rehydration buffer. The data show that such an increase in the amount of α -crystallin does not result in higher cell survival after rehydration. The conclusion from these data is that α -crystallin improves the survival of freeze-dried and rehydrated erythrocytic cells, as assessed by the decrease in hemolysis from 68% (in cells that have not been loaded in the presence of α -crystallin) to 49% (in cells loaded in the presence of α -crystallin).

It has been further discovered that when Zn^{2+} ions are added to the rehydration buffer, there is a decrease in the percent hemolysis of rehydrated erythrocytic cells, suggesting that these ions have beneficial effect on cell survival after freeze-drying. Zn^{2+} ions stabilize thermally labile enzymes during drying. Rehydration experiments were performed combining α -

crystallin and Zn^{2+} ions, and applying 5 min pre-hydration. Under these conditions, 62% of the cells survived the rehydration step, indicating that the beneficial effect of these treatments is additive. Figure 22 shows the combined effect of α -crystallin, Zn^{2+} and pre-hydration on the survival of erythrocytic cells. When the cells were loaded in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate (Figure 22, data set labeled as **T**), freeze-dried in 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HSA and directly rehydrated at 37°C for 10 min in 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA, the hemolysis was 63%. When the cells were loaded in a buffer containing 800 mM trehalose, 100 mOsm ADSOL, 6.6 mM Na-phosphate and α -crystallin (1.2 mg/ml), freeze-dried in 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HAS, pre-hydrated at 37°C for 5 min, and then fully rehydrated at 37°C for 10 min in 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA, the hemolysis was 47% (Figure 22, data set labeled as **[T+ α C]+5min**) . When cells were loaded in a buffer containing 1.2 mg/ml α -crystallin, 800 mM trehalose, 100 mOsm ADSOL, 6.6 mM Na-phosphate, freeze-dried in 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HSA, pre-hydrated for 5 min at 37°C, and rehydrated at 37°C for 10 min in a buffer containing 500 μM ZnSO_4 , 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA, the hemolysis was only 38% (Figure 22, data set labeled as **[T+ α C]+5min+Zn**), giving rise to 62% cell survival.

The levels of ATP and 2,3-DPG were followed during rehydration of freeze-dried erythrocytic cells. Incubation of the rehydrated cells in a buffer supplemented with rejuvenation solution led to considerable increase in the ATP and 2,3-DPG synthesis. Figure 23 shows the levels of the two metabolites during 10 min and 60 min incubation at 37°C in a rehydration

buffer containing 141 mM trehalose, 15% HES, 2.5% HSA and the following rejuvenation supplements: 100 mM pyruvate, 100 mM inosine, 100 mM Na-phosphate and 5 mM adenine. The rejuvenation solution is referred as PIPA. Cells were loaded in 800 mM trehalose, 100 mOsm ADSOL and 6.6 mM Na-phosphate at 38-41°C for 6 hours. They were freeze-dried in 250 mM trehalose, 20 mOsm ADSOL, 15% HES and 2.5% HSA, and rehydrated at 37°C for 10 min in 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA (Figure 23, data set labeled as **141**). Without applying rejuvenation solution, the levels of both ATP and 2,3-DPG are low. However, when such cells were rehydrated in buffer containing 141 mM trehalose, 75 mOsm PBS, 11.25% HES and 1.875% HSA, supplemented with PIPA, the levels of ATP and 2,3-DPG were increased (Figure 23, data set labeled as **141PIPA**). These results show that supplementation of the rehydration medium with 100 mM pyruvate, 100 mM inosine, 100 mM Na-phosphate and 5 mM adenine increases the synthesis of these two vital metabolites and can be applied during reconstitution of freeze-dried erythrocytic cells.

Embodiments of the present invention will be illustrated by the following set forth examples which are being given to set forth the presently known best mode and by way of illustration only and not by way of any limitation. It is to be understood that all materials, chemical compositions and procedures referred to below, but not explained, are well documented in published literature and known to those artisans possessing skill in the art. All materials and chemical compositions whose source(s) are not stated below are readily available from commercial suppliers, who are also known to those artisans possessing skill in the art. All parameters such as concentrations, mixing proportions, temperatures, rates, compounds, etc., submitted in these examples are not to be

construed to unduly limit the scope of the invention.

Abbreviations used in the examples, and elsewhere, are as follows:

DMSO = dimethylsulfoxide

ADP = adenosine diphosphate

PGE1 = prostaglandin E1

HES = hydroxy ethyl starch

FTIR = Fourier transform infrared spectroscopy

EGTA = ethylene glycol-bis(2-aminoethyl ether) N,N,N',N',
tetra-acetic acid

TES = N-tris (hydroxymethyl) methyl-2-aminoethane-sulfonic
acid

HEPES = N-(2-hydroxyl ethyl) piperazine-N'-(2-
ethanesulfonic acid)

PBS = phosphate buffered saline

HSA = human serum albumin

BSA = bovine serum albumin

ACD = citric acid, citrate, and dextrose

M β CD = methyl- β -cyclodextrin

EXAMPLE 1

Washing of Platelets. Platelet concentrations were obtained from the Sacramento blood center or from volunteers in our laboratory. Platelet rich plasma was centrifuged for 8 minutes at 320 x g to remove erythrocytes and leukocytes. The supernatant was pelleted and washed two times (480 x g for 22

minutes, 480 x g for 15 minutes) in buffer A (100 mM NaCl, 10 mM KCl, 10 mM EGTA, 10 mM imidazole, pH 6.8). Platelet counts were obtained on a Coulter counter T890 (Coulter, Inc., Miami, Florida).

Loading of Lucifer Yellow CH into Platelets. A fluorescent dye, lucifer yellow CH (LYCH), was used as a marker for penetration of the membrane by a solute. Washed platelets in a concentration of $1-2 \times 10^9$ platelets/ml were incubated at various temperatures in the presence of 1-20 mg/ml LYCH. Incubation temperatures and incubation times were chosen as indicated. After incubation the platelets suspensions were spun down for 20 s at 14,000 RPM (table centrifuge), resuspended in buffer A, spun down for 20 s in buffer A and resuspended. Platelet counts were obtained on a Coulter counter and the samples were pelleted (centrifugation for 45 s at 14,000 RPM, table centrifuge). The pellet was lysed in 0.1% Triton buffer (10 mM TES, 50 mM KCl, pH 6.8). The fluorescence of the lysate was measured on a Perkin-Elmer LSS spectrofluorimeter with excitation at 428 nm (SW 10 nm) and emission at 530 nm (SW 10 nm). Uptake was calculated for each sample as nanograms of LYCH per cell using a standard curve of LYCH in lysate buffer. Standard curves of LYCH, were found to be linear up to 2000 ng ml^{-1} .

Visualization of cell-associated Lucifer Yellow. LYCH loaded platelets were viewed on a fluorescence microscope (Zeiss) employing a fluorescein filter set for fluorescence microscopy. Platelets were studied either directly after incubation or after fixation with 1% paraformaldehyde in buffer. Fixed cells were settled on poly-L-lysine coated cover slides and mounted in glycerol.

Loading of Platelets with Trehalose. Washed platelets in a concentration of $1-2 \times 10^9$ platelets/ml were incubated at various

temperatures in the presence of 1-20 mg/ml trehalose. Incubation temperatures were chosen from 4°C to 37°C. Incubation times were varied from 0.5 to 4 hours. After incubation the platelet solutions were washed in buffer A two times (by centrifugation at 14,000 RPM for 20 s in a table centrifuge). Platelet counts were obtained on a coulter counter. Platelets were pelleted (45 S at 14,000 RPM) and sugars were extracted from the pellet using 80% methanol. The samples were heated for 30 minutes at 80°C. The methanol was 10 evaporated with nitrogen, and the samples were kept dry and redissolved in H₂O prior to analysis. The amount of trehalose in the platelets was quantified using the anthrone reaction (Umbreit et al., *Mamometric and Biochemical Techniques*, 5th Edition, 1972). Samples were redissolved in 3 ml H₂O and 6 ml anthrone reagents (2 g anthrone dissolved in 10M sulfuric acid). After vortex mixing, the samples were placed in a boiling water bath for 3 minutes. Then the samples were cooled on ice and the absorbance was measured at 620 nm on a Perkin Elmer spectrophotometer. The amount of platelet associated trehalose was determined using a standard curve of trehalose. Standard curves of trehalose were found to be linear from 6 to 300 µg trehalose per test tube.

Quantification of Trehalose and LYCH Concentration. Uptake was calculated for each sample as micrograms of trehalose or LYCH per platelet. The internal trehalose concentration was calculated assuming a platelet radius of 1.2 µm and by assuming that 50% of the platelet volume is taken up by the cytosol (rest is membranes). The loading efficiency was determined from the cytosolic trehalose or LYCH concentration and the concentration in the loading buffer.

Fig. 1 shows the effect of temperature on the loading efficiency of trehalose into human platelets after a 4 hour

incubation period with 50 mM external trehalose. The effect of the temperature on the trehalose uptake showed a similar trend as the LYCH uptake. The trehalose uptake is relatively low at temperatures of 22°C and below (below 5%), but at 37°C the loading efficiency of trehalose is 35% after 4 hours.

When the time course of trehalose uptake is studied at 37°C, a biphasic curve can be seen (Fig. 2). The trehalose uptake is initially slow (2.8×10^{-11} mol/m²s from 0 to 2 hours), but after 2 hours a rapid linear uptake of 3.3×10^{-10} mol/m²s can be observed. The loading efficiency increases up to 61% after an incubation period of 4 hours. This high loading efficiency is a strong indication that the trehalose is homogeneously distributed in the platelets rather than located in pinocytosed vesicles.

The uptake of trehalose as a function of the external trehalose concentration is shown in Fig. 3, which graphically illustrates the internal trehalose concentration of human platelets versus external trehalose concentration as a function of temperature at a constant incubation or loading time. The uptake of trehalose is linear in the range from 0 to 30 mM external trehalose. The highest internal trehalose concentration is obtained with 50 mM external trehalose. At higher concentrations than 50 mM the internal trehalose concentration decreases again. Even when the loading buffer at these high trehalose concentrations is corrected for isotonicity by adjusting the salt concentration, the loading efficiency remains low. Platelets become swollen after 4 hours incubation in 75 mM trehalose. Figure 4 graphically illustrates the loading efficiency of trehalose into human platelets as a function of external trehalose concentration.

The stability of the platelets during a 4 hours incubation period was studied using microscopy and flow cytometric analysis. No morphological changes were observed after 4 hours incubation of platelets at 37°C in the presence of 25 mM external trehalose. Flow cytometric analysis of the platelets showed that the platelet population is very stable during 4 hours incubation. No signs of microvesicle formation could be observed after 4 hours incubation, as can be judged by the stable relative proportion of microvesicle gated cells (less than 3%). The formation of microvesicles is usually considered as the first sign of platelet activation (Owens et al., *Trans. Med. Rev.*, 8, 27-44, 1994). Characteristic antigens of platelet activation include: glycoprotein 53 (gp53, a lysosomal membrane marker), PECAM-1 (platelet endothelial cell adhesion molecule-1, an alpha granule constituent), and P-selection (an alpha granule membrane protein).

EXAMPLE 2

Figure 5 graphically illustrates the loading efficiency of trehalose into human erythrocytic cells as a function of external trehalose concentration at respective temperatures of 4°C and 37°C. Erythrocytic cells were exposed to trehalose for 18 hours at either 4°C or 37°C. The trehalose concentration in the incubation medium varied between 230 mM and 1000 mM. Each incubation buffer contained trehalose (between 230 mM and 1000 mM) and 100 mOsm PBS pH 7.2. Increase in the trehalose concentration in the loading medium results in an increase in the sugar uptake, reaching about 100 mM cytoplasmic trehalose in erythrocytes incubated in 1000 mM trehalose and 100

mOsm PBS. At 4° C, the uptake was very limited, being about 25 mM. The trehalose intake was measured using anthrone assay and confirmed by high performance liquid chromatography. It is clear that there was substantial loading at 37° C, but not at 4° C. Furthermore, trehalose loading was not significant unless the extracellular cellular trehalose concentration gave a hyperosmotic pressure. Since intracellular osmolarity for erythrocytic cells is about 300 mOsm, it is clear that raising the extracellular osmolarity was required for more effective loading of trehalose.

EXAMPLE 3

Figure 6 graphically illustrates the fragility index of erythrocytic cells incubated overnight at respective temperatures of 4° C and 37° C in the presence of and as a function of increasing intracellular trehalose concentrations. The osmotic fragility index was generated by the extent of hemolysis as a function of the NaCl concentration. The erythrocytic cells that had been loaded in trehalose solutions (between 250 mM and 1000 mM) in 100 mOsm PBS were suspended in increasing concentrations of NaCl (between 50 and 600 mOsm NaCl). The percent hemolysis measured after resuspending the loaded cells in NaCl represents the fragility index. The data show that the erythrocytic cells were stable osmotically in trehalose media with concentrations between 250 mM and 800 mM trehalose at both 37° C and 4° C. In 1000 mM trehalose at 37° C, there is a high increase in the fragility index suggesting that the cells were unstable in this medium (1000 mM trehalose in 100 mOsm PBS).

EXAMPLE 4

Figure 7 graphically illustrates trehalose uptake (i.e., intracellular trehalose mM) and hemolysis (i.e., % hemolysis) as a function of incubation temperature ($^{\circ}\text{C}$). The incubation temperature was varied between 4°C and 37°C . The erythrocytic cells were incubated for 6 hours in 800 mM trehalose in 100 mOsm PBS pH 7.2. Between 4°C and 30°C , the cytoplasmic trehalose was very low (between 1 and 4 mM). It was considerably increased (up to 35 mM cytoplasmic trehalose) during 6 hours incubation at 37°C .

EXAMPLE 5

Figure 8 graphically illustrates intracellular trehalose concentration (mM) as a function of the osmolarity of the washing buffer. Earlier morphological data showed that along with discoid erythrocytic cells, there is about 20% of cells with modified shape (spherocytes and schistocytes). The issue was what was the loading capacity of these cells and how much they contribute to the amount of trehalose that was to be detected. This issue was investigated by washing the trehalose loaded erythrocytic cells (loaded at 35°C for 16 hours in 800 mM trehalose in 100 mOsm PBS pH 7.2) in buffers with different osmolarity (300 mOsm PBS or 900 mOsm PBS) and estimating the cytoplasmic sugar concentration. The loaded cells were washed with either 300 mOsm PBS pH 7.2 (which is the isotonic medium for erythrocytic cells) or 900 mOsm PBS pH 7.2 (which matches the tonicity of the loading medium). The data in Figure 8

illustrated that there is a decrease in the intracellular sugar concentration suggesting that a fraction of the cells was lost during the washing procedure.

EXAMPLE 6

Figure 9 graphically illustrates % hemolysis of loaded cells as a function of time (hours) of incubation in 300 mOsm PBS pH 7.2. The erythrocytic cells were loaded in 700 mM trehalose in 100 mOsm PBS pH 7.2 at 35° C for 16 hours. After the loading step, the cells were incubated in 300 mOsm PBS pH 7.2 between 1 and 25 hours. This buffer was used for washing the loaded cells and it was important to determine the stability of the loaded cells when they were suspended in this buffer. Cell stability was assessed by measuring the percent hemolysis over the course of 25 hours. The data show an increase in the percent hemolysis in the first 7 hours (from about 7 % to about 25 % hemolysis). Between 7 and 25 hours, there was only a small change in the percent hemolysis (from about 25 % to about 30 %).

EXAMPLE 7

Figure 10 graphically illustrates time course (incubation time, hours) of hemolysis(%) of trehalose- loaded cells as a function of the composition of the incubation buffer. Erythrocytic cells were loaded in 700 mM trehalose in 100 mOsm PBS pH 7.2 for 16 hours at 35° C. They were washed with 300 mOsm PBS pH 7.2 and were subsequently transferred to storage media

with increasing trehalose concentrations (between 100 mM and 300 mM trehalose) added to 100 mOsm PBS pH 7.2. The percent hemolysis was measured over the course of 3 hours. The results show that the percent hemolysis is reduced with increase of trehalose concentration (from about 25 % in 300 mOsm PBS with no trehalose, to about 4 % in 300 mM trehalose in 100 mOsm PBS).

EXAMPLE 8

Figure 11 graphically illustrates time course (incubation time, hours) of hemolysis(%) of trehalose- loaded cells as a function of the composition of the incubation buffer, and illustrating that HES and albumin (HSA) do not have any detrimental effect on cell hemolysis during incubation. We tested the effect of 15 % HES and 30 % HES, and the effect of 2.5 % albumin (HSA) and 5 % albumin, as well as a combination of 2.5 % albumin and 15 % HES. The results show that a combination of the two compounds provides the lowest percent hemolysis of trehalose-loaded erythrocytic cells.

EXAMPLE 9

Figure 12 is a 40X picture of rehydrated erythrocytic cells having no intracellular trehalose prior to freeze-drying. These cells had not been loaded with trehalose, and were freeze-dried in 300 mM trehalose in 100 mOsm PBS, 15 % HES and 2.5 % albumin. Rehydration was done in 188 mM trehalose, 100 mOsm PBS, 15 % HES and 2.5 % albumin. Figure 12 illustrates that when the cells are not loaded with trehalose, and are freeze-dried and rehydrated,

there are mostly cells with modified shape (spherocytes) and cell debris.

Figure 13 is a 40X picture of rehydrated erythrocytic cells having 3 mM intracellular trehalose, after initially trehalose-loading the cells in 400 mM trehalose in 100 mOsm PBS at 35° C for 16 hours and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Rehydration was done in 188 mM trehalose, 100 mOsm PBS, 15 % HES and 2.5 % albumin. Figure 13 displays erythrocytic cells containing 3 mM cytoplasmic trehalose, freeze-dried and rehydrated. Cells with biconcave shape were seen, as well as cells with modified morphology, suggesting that trehalose loading of erythrocytic cells provide higher number of cells with preserved morphology.

Figure 14 is a 40X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading in 800 mM trehalose in 100 mOsm PBS at 35° C for 16 hours and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Rehydration was done in 188 mM trehalose, 100 mOsm PBS, 15 % HES and 2.5 % albumin. Figure 14 shows erythrocytic cells containing 60 mM cytoplasmic trehalose, freeze-dried and rehydrated. The typical biconcave shape and cell integrity appear well preserved.

Figure 15 is a 100X picture of rehydrated erythrocytic cells having 60 mM intracellular trehalose, after initially trehalose-loading in 800 mM trehalose in 100 mOsm PBS at 35° C for 16 hours and freeze-drying the cells in 300 mM trehalose/100 mOsm PBS, 15% HES and 2.5% HSA. Rehydration was done in 188 mM trehalose, 100 mOsm PBS, 15 % HES and 2.5 % albumin. Figure 15 shows erythrocytic cells containing 60 mM cytoplasmic trehalose, freeze-dried and rehydrated at higher

magnification (100X) confirming that the cell shape is mostly biconcave and cell integrity is preserved.

EXAMPLE 10

Figure 16 is a graph of hemolysis(%) of trehalose loaded, freeze-dried and rehydrated erythrocytic cells as a function of intracellular trehalose concentration (mM), graphically illustrating the effect of cytoplasmic trehalose on the survival of rehydrated erythrocytic cells. The cells were loaded in media with different trehalose concentration (400, 500, 600, 700 and 800 mM trehalose) in 100 mOsm PBS pH 7.2. The cells were freeze-dried in 300 mM trehalose in 100 mOsm PBS, 15 % HES and 2.54 % HSA and rehydrated in a buffer containing 188 mM trehalose, 15 % HES and 2.5 % HSA. Figure 16 shows that higher extent of trehalose loading of erythrocytic cells confers higher percent of survival, assessed by the percent hemolysis of the rehydrated cells.

EXAMPLE 11

Figure 17 is a graph of mean corpuscular hemoglobin of trehalose-loaded, freeze-dried and rehydrated erythrocytic cells as a function of intracellular trehalose concentration (mM), graphically illustrating that as the concentration of intracellular trehalose increases for rehydrated erythrocytic cells, the mean corpuscular hemoglobin (the amount of hemoglobin found in intact erythrocytic cells) also increases for rehydrated erythrocytic cells. The cells had been loaded in

media with different trehalose concentration (400 mM trehalose, 500 mM trehalose, 600 mM trehalose, 700 mM trehalose and 800 mM trehalose) in 100 mOsm PBS pH 7.2. The cells were freeze-dried in 300 mM trehalose in 100 mOsm PBS, 15 % HES and 2.54 % HSA and rehydrated in a buffer containing 188 mM trehalose, 15 % HES and 2.5 % HSA. The data demonstrate that higher cytoplasmic trehalose increases the amount of mean corpuscular hemoglobin in rehydrated cells (from about 7 pg in cells loaded in 400 mM trehalose to about 14 pg in cells loaded in 800 mM trehalose).

EXAMPLE 12

The following protocol provided significant survival of freeze-dried and rehydrated erythrocytic cells. Loading buffer comprised 800 mM trehalose in a salt solution of 100 mOsm PBS. The incubation time was 16 hours at a temperature of about 35° C. After the cells were loaded, they were subsequently washed in a washing buffer comprising 300 mM in a salt solution of 100 mOsm PBS. Within 3 hours after washing the loaded cells, the wash loaded cells were freeze-dried in freeze-drying buffer comprising about 300 mM trehalose, about 100 mOsm PBS, about 2.5 % by wt. HSA, and about 15 % by wt. HES. After the freeze-drying procedure the cells had about 75 mM trehalose, about 25 mOsm PBS, about 0.6 % by wt. HSA and about 4.0 % by wt. HES left in the cells. The freeze-dried cells were then reconstituted at about 37° C for about 10 minutes in a rehydration buffer comprising about 188 mM trehalose, about 100 mOsm PBS, about 2.5 % by wt. HSA and about 15.0 % by wt. HES. After rehydration, less than about 5 % of the cells lysed.

Conclusion

Embodiments of the present invention provide that trehalose, a sugar found at high concentrations in organisms that normally survive dehydration, may be used to preserve biological structures in the dry state. Cells may be loaded with trehalose under the previously specified conditions, and the loaded cells can be freeze dried with excellent recovery.

While the present invention has been described herein with reference to particular embodiments thereof, a latitude of modification, various changes and substitutions are intended in the foregoing disclosure, and it will be appreciated that in some instances some features of the invention will be employed without a corresponding use of other features without departing from the scope and spirit of the invention as set forth. Therefore, many modifications may be made to adapt a particular situation or material to the teachings of the invention without departing from the essential scope and spirit of the present invention. It is intended that the invention not be limited to the particular embodiment disclosed as the best mode contemplated for carrying out this invention, but that the invention will include all embodiments and equivalents falling within the scope of the appended claims.

What Is Claimed Is:

1. A method for loading a solute into a cell comprising:
disposing a cell in a solution having a solute concentration having a temperature of at least 25° C and of sufficient magnitude to produce hyperosmotic pressure on the cell for transferring a solute from the solution into the cell.
2. The method of Claim 1 wherein said solute concentration includes an extracellular cellular solute concentration for elevating extracellular osmolarity within the solution to a value which is greater than a value of the intracellular osmolarity of the cell.
3. The method of Claim 1 wherein said transferring a solute is by fluid phase endocytosis.
4. The method of Claim 1 wherein said solute comprises trehalose and said cell comprises an erythrocytic cell.
5. The method of Claim 4 wherein said transferring of trehalose from the solution into the erythrocytic cell is without degradation of the trehalose.
6. The method of Claim 4 wherein a gradient of trehalose concentration within the erythrocytic cell to extracellular trehalose concentration within the solution ranges from about 0.130 to about 0.200.

7. The method of Claim 4 wherein said solute solution has a trehalose concentration ranging from about 320 mM to about 4000 mM.

8. The method of Claim 1 additionally comprising

preventing a decrease in a loading efficiency gradient in the loading of the solute into the cell.

9. The method of Claim 8 wherein said solute comprises an oligosaccharide and said preventing a decrease in a loading efficiency gradient in the loading of the oligosaccharide into the cell comprises maintaining a concentration of the oligosaccharide in the oligosaccharide solution below a concentration ranging from about 35 mM to about 65 mM.

10. The method of Claim 8 wherein said solute comprises an oligosaccharide and said preventing a decrease in a loading efficiency gradient in the loading of the oligosaccharide into the cell comprises maintaining a positive gradient of loading efficiency to concentration of the oligosaccharide in the oligosaccharide solution.

11. A cell produced in accordance with the method of Claim 1.

12. A method for loading trehalose into an erythrocytic cell comprising disposing an erythrocytic cell in a trehalose solution having a temperature of at least 25° C and a trehalose concentration of at least about 25 % greater than the intracellular osmolarity of the erythrocytic cell for loading the trehalose into the erythrocytic cell.

13. The method of Claim 12 wherein said loading the trehalose into the erythrocytic cell is by fluid phase endocytosis.
14. The method of Claim 12 wherein said loading of the trehalose from the trehalose solution into the erythrocytic cell is without degradation of the trehalose.
15. The method of Claim 12 said loading of the trehalose produces a loaded erythrocytic cell having a gradient of loaded trehalose concentration within the erythrocytic cell to extracellular trehalose concentration within the trehalose solution ranging from about 0.130 to about 0.200.
16. The method of Claim 12 wherein said trehalose solution has a trehalose concentration ranging from about 25 % to at least about 1000% greater than the intracellular osmolarity of the erythrocytic cell.
17. An erythrocytic cell produced in accordance with the method of Claim 12.
18. A method for preparing a dehydrated composition comprising:
loading cells in a loading solution having a salt solution and a solute for producing loaded cells; and
lyophilizing the loaded cells in a freeze-drying solution having a drying salt solution, the solute, an inert substance and a protein to produce a dehydrated composition.
19. The method of Claim 18 wherein said loading solution comprises at least about 200 mM of the solute and at least about 75 mOsm of the salt solution.

20. The method of Claim 19 wherein said freeze-drying solution comprises at least about 50 mM of the solute; at least about 2.0 % by weight of the inert substance; at least about 0.5 % by weight of the protein, and at least about 25 mOsm for an osmolarity of the salt solution.

21. A method for reducing hemolysis in treating cells comprising loading cells in an incubation period with a loading solution comprising a salt solution and at least about 200 mM of a solute to reduce hemolysis of the cells to less than about 10 %.

22. The method of Claim 21 wherein said loading solution additionally comprises a starch and a protein, and said hemolysis of the cells is reduced to less than about 5 %.

23. The method of Claim 21 additionally comprising washing the loaded cells, and drying washed cells within 2 hours after washing to assist in maintaining hemolysis below about 10 %.

24. A method for stabilizing cells comprising loading cells in a loading solution to produce cells having an effective amount of a solute for possessing a mean corpuscular hemoglobin (pg) greater than about 10.

25. The method of Claim 24 wherein said mean corpuscular hemoglobin (pg) is greater than about 14.

26. The method of Claim 24 wherein said effective amount of a solute is greater than about 50 mM.

27. A method for reconstituting dried cells comprising

drying solute-loaded cells in a drying solution having a salt solution, a solute, an inert substance, and a protein to produce dried cells; and

reconstituting the dried cells in a rehydration solution having the salt solution, the solute, the inert substance, and the protein to produce reconstituted cells.

28. The method of Claim 27 wherein said drying solution comprises at least about 50 mM of the solute; at least about 25 mOsm osmolarity of the salt solution; at least about 2.0 % by weight of the inert substance; and at least about 0.5 % by weight of the protein.

29. The method of Claim 28 wherein said rehydration solution comprises at least about 50 mM of the solute; at least about 25 mOsm osmolarity of the salt solution; at least about 2.0 % by weight of the inert substance; and at least about 0.5 % by weight of the protein.

30. The method of Claim 27 wherein said dried cells comprise from about 25 mM to about 300 mM of the solute; from about 5 mOsm to about 100 mOsm osmolarity for the salt solution; from about 0.1 % by weight to about 2.5 % by weight of the protein; and from about 1.0 % by weight to about 15.0 % by weight of the inert substance.

31. The method of Claim 30 wherein solute comprises trehalose, said salt solution comprises PBS, said protein comprises albumin, and said inert substance comprises a starch.

32. The method of Claim 31 wherein said dried cells comprise from about 60 mM to about 80 mM trehalose, from about 10 mOsm to about 40 mOsm PBS, from about 0.3 % by weight to about 9.0 % by weight albumin, and about 1.0 % by weight to about 4.0 % by weight starch.

33. The method of Claim 21 additionally comprising adding an inert substance and/or a protein to the loading solution to further reduce hemolysis.

CELLS AND METHOD FOR PRESERVING CELLS

Abstract of the Disclosure

A dehydrated composition is provided that includes freeze-dried cells. A method for loading a solute into a cell comprising disposing a cell in a solution having at temperature of at least 25° C and a solute concentration of sufficient magnitude to produce hyperosmotic pressure on the cell for transferring a solute from the solution into the cell. A method for reconstituting dried cells by drying loaded cells in a drying solution and reconstituting dried cells in a rehydration solution. A method for stabilizing cells by producing cells having an effective amount of a solute and a mean corpuscular hemoglobin greater than about 10.

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




FIG. 1




FIG. 2

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




FIG. 3




FIG. 4

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




Fig. 5

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US

**Fragility index of RBCs incubated overnight at 4 or 37°C
in the presence of increasing trehalose concentrations**




Figure 6

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




Fig 7

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US

Intracellular trehalose concentration as a function of the osmolarity of the washing buffer.




Figure 8

Percent hemolysis of trehalose loaded RBCs as a function of time of incubation in 300 mOsm PBS. RBCs were loaded in 700 mM trehalose/100 mOsm PBS at 35°C for 16 hours and were incubated in 300 mOsm PBS.




Figure 9

Percent hemolysis of trehalose loaded RBCs as a function of the composition of the incubation buffer. RBCs were loaded in 700 mM trehalose/100 mOsm PBS at 35°C for 16 hours




Figure 10

Percent hemolysis of trehalose loaded RBCs as a function of the composition of the incubation buffer.
 RBCs were loaded in 700 mM trehalose/100 mOsm PBS at 35°C for 16 hours




Figure 11

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US



Fig 12

0 mM

40X

BEST AVAILABLE COPY

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




Fig 13

3 mM

CELLS AND METHOD FOR PRESERVING CELLS


Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US

60 mM

40X

Fig 14



CELLS AND METHOD FOR PRESERVING CELLS
Inventors: John H. Crowe et al.
Atty Docket No. 010023-000800US




Fig 15

50 mM

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




Fig 16


Intracellular trehalose (mM)

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US

Fig 17



ATP ($\mu\text{mol/g Hb}$) in erythrocytes incubated in 800 mM trehalose and different buffers
(see legend) as a function of time of incubation at 40°C




Figure 1

2,3-DPG level in erythrocytes incubated in different buffers as a function of time of incubation at 40°C





Figure 19

Effect of time of prehydration on the survival of freeze-dried
and rehydrated erythrocytes



Effect of α -crystallin on the survival of freeze-dried and rehydrated erythrocytes

CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US




Figure 21

Effect of pre-hydration (5 min), α -crystallin (1.2 mg/ml) and Zn^{2+} (500 μl) on the survival of freeze-dried and rehydrated erythrocytes.



CELLS AND METHOD FOR PRESERVING CELLS

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000800US

Effect of rejuvenating buffer on the synthesis of ATP and 2,3-DPG in rehydrated erythrocytes.

CELLS AND METHOD FOR PRESERVING CELLS
Inventors: John H. Crowe et al.
Atty Docket No. 010023-000800US




Figure 23

PATENT

Applicant: John H. Crowe et al.
Serial No.: Unknown
Filed: August 6, 2003
Title: CELLS AND METHOD FOR PRESERVING CELLS

Group: Unknown
Examiner: Unknown
Docket No. 010023-000800

COMBINED DECLARATION AND POWER OF ATTORNEY
(ORIGINAL, DESIGN, SUPPLEMENTAL, NATIONAL STAGE OF PCT, DIVISIONAL, CONTINUATION OR
CONTINUATION-IN-PART)

As a below named inventor, I hereby declare that:

TYPE OF DECLARATION

This declaration is of the following type:

- ☒ original (regular)
- ☐ design
- ☐ supplemental-continuation of PCT Application in the U.S.

NOTE: If the declaration is for an International Application being filed as a divisional; continuation, or continuation-in-part, do not check the next item; check appropriate one of last three items.

- ☐ national stage of PCT divisional continuation continuation-in-part

INVENTORSHIP IDENTIFICATION

WARNING: If the inventors are each not the inventors of all of the claims, an explanation of the facts, including the ownership of all the claims at the time the last claimed invention was made, should be submitted.

My residence, post office address, and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention described in

SPECIFICATION IDENTIFICATION

- ☒ the specification filed herewith by the above-named inventors, with the title listed above.
- ☐ the specification filed herewith by the above-named inventors, with the title listed above, and which was amended by the Preliminary Amendment filed herewith.
- ☐ the specification identified above, as amended by the Preliminary Amendment filed herewith.
- ☐ the specification identified above, Serial No. _____ filed on _____.
- ☐ PCT International Application No. _____ filed on _____, and amended under PCT Article 19 on _____ (if applicable).

ACKNOWLEDGMENT OF REVIEW OF PAPERS AND DUTY OF CANDOR

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. §1.56.

PRIORITY CLAIM (35 U.S.C. § 119(a)-(d))

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119(a)-(d) of any foreign application(s) for patent or inventor's certificate or under 35 U.S.C. § 365(b) of any PCT international application(s) designating at least one country other than the United States of America listed below, and have also identified below any foreign application(s) or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

☐ No such applications have been filed.

☐ Such applications have been filed as follows:

PRIOR FOREIGN/PCT APPLICATION(S) FILED WITHIN 12 MONTHS (6 MONTHS FOR DESIGN) PRIOR TO THIS APPLICATION AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. § 119(a)-(d)

COUNTRY (or indicate if PCT)	APPLICATION NUMBER	FILING DATE	PRIORITY CLAIMED UNDER 37 U.S.C. § 119	
			<input type="checkbox"/> YES	<input type="checkbox"/> NO
			<input type="checkbox"/> YES	<input type="checkbox"/> NO
			<input type="checkbox"/> YES	<input type="checkbox"/> NO
			<input type="checkbox"/> YES	<input type="checkbox"/> NO

CLAIM FOR BENEFIT OF PRIOR U.S. PROVISIONAL APPLICATION(S) (35 U.S.C. § 119(e))

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional applications(s) listed below:

PROVISIONAL APPLICATION SERIAL NUMBER	FILING DATE OF PROVISIONAL APPLICATION

CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S) UNDER 35 U.S.C. § 120

(Complete this part only if this is a divisional, continuation, CIP or national stage of PCT.)

I/We hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability under 37 C.F.R. § 1.56, which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.

PRIOR U.S. PROVISIONAL APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. § 120:

1. U.S. Applications:

SERIAL NUMBER	FILING DATE

2. PCT Applications Designating The U.S.

PCT APPLICATION NUMBER	PCT FILING DATE	U.S. SERIAL NUMBER

POWER OF ATTORNEY

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the United States Patent and Trademark Office connected therewith:

John W. Carpenter	26,447
Charles J. Kulas	35,809

☐ Attached as part of this declaration and power of attorney is the authorization of the above-named attorney to accept and follow instructions from my representative(s).

SEND CORRESPONDENCE TO:

Carpenter & Kulas, LLP
 1900 Embarcadero Road, Suite 109
 Palo Alto, CA 94303

DIRECT TELEPHONE CALLS TO:

John W. Carpenter
 (650) 842-0303

DECLARATION

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code,, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

SIGNATURE(S)

Full name of first inventor: John H. Crowe

Inventor's
 Signature: _____

Date: _____

Country of Citizenship: USA

Residence: 1111 Cottonwood Court, Davis, CA 95616

P.O. Address: same

Full name of second inventor: Fern Tablin

Inventor's

Signature: _____

Date: _____

Country of Citizenship: USA

Residence: 608 Huble Street, Davis, CA 95616

P.O. Address: same

Full name of third inventor: Nelly M. Tsvetkova

Inventor's

Signature: _____

Date: _____

Country of Citizenship: Bulgarian

Residence: 3114 Newport Terrace, Davis, CA 95616

P.O. Address: same

Full name of fourth inventor: Zsolt Torok

Inventor's

Signature: _____

Date: _____

Country of Citizenship: Hungary

Residence: 4501 Alhambra Drive, #228, Davis, CA 95616

P.O. Address: same

Full name of fifth inventor: Rachna Bali

Inventor's

Signature: _____

Date: _____

Country of Citizenship: USA

Residence: 1691 Columbus Road, West Sacramento, CA 95691

P.O. Address: same

Full name of sixth inventor: Gyana R. Satpathy

Inventor's

Signature: _____

Date: _____

Country of Citizenship: India

Residence: 424 Russell Park #2, Davis, CA 95616

P.O. Address: same

Full name of seventh inventor: Denis M. Dwyre

Inventor's

Signature: _____

Date: _____

Country of Citizenship: USA

Residence: 134 Ashford Place, Iowa City, IA 52245

P.O. Address: same

CHECK PROPER BOX(ES) FOR ANY OF THE FOLLOWING ADDED PAGE(S) WHICH FORM A PART OF THIS DECLARATION

- ☒ Signature for third and subsequent joint inventors.
Number of pages added 1
- ☐ Signature by administrator(trix), executor(trix) or legal representative for deceased or incapacitated inventor.
Number of pages added _____
- ☐ Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR 1.47.
Number of pages added _____
- ☐ Signature by one joint inventor on behalf of deceased inventor(s) where legal representative cannot be appointed in time under 37 CFR 1.47.
Number of pages added _____
- ☐ Authorization of attorney(s) to accept and follow instructions from representative.

☒ This Declaration ends with this page.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/025395

International filing date: 06 August 2004 (06.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 10/635,795
Filing date: 06 August 2003 (06.08.2003)

Date of receipt at the International Bureau: 13 December 2004 (13.12.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.